

# Numerical Solutions to Differential Equations

## Lecture Notes #4 — Stability Regions Revisited & Runge-Kutta Methods

Peter Blomgren,  
(blomgren.peter@gmail.com)

Department of Mathematics and Statistics  
Dynamical Systems Group  
Computational Sciences Research Center  
San Diego State University  
San Diego, CA 92182-7720  
<http://terminus.sdsu.edu/>

Spring 2015

Stability Regions Revisited & RK-Methods — (1/34)

## Outline

- 1 Recap: Last Lecture
- 2 Finding Stability Regions
  - Euler's Method
  - Taylor Series Methods
- 3 Runge-Kutta Methods
  - Introduction
  - s-stage RK-methods
  - Types of RK-methods
  - Derivation

Stability Regions Revisited & RK-Methods — (2/34)

## Last Lecture: Quick Review

### Euler's Method:

- Analysis: Local Truncation Error (LTE), Consistency, Accuracy, Stability (Region of Stability), Convergence

### Improvements:

- Higher order Taylor Series Methods
- Multi-Point Methods
  - Heun's Method
  - Euler's "Midpoint Method"

Stability Regions Revisited & RK-Methods — (3/34)

## Stability Regions Revisited

Recall: Euler's Method

$$y_{n+1} = y_n + hf(t_n, y_n), \quad y(t_0) = y_0$$

applied to

$$y'(t) = \lambda y(t)$$

gives

$$y_{n+1} = y_n + h\lambda y_n = (1 + h\lambda)y_n = (1 + h\lambda)^{n+1}y_0.$$

The stability criterion is (non-exponential growth):

$$|1 + h\lambda| \leq 1$$

Stability Regions Revisited & RK-Methods — (4/34)


## Finding the Stability Region

How do we find the stability region from the expression

$$|1 + h\lambda| \leq 1.$$

The boundary of the region is given by

$$1 + h\lambda = e^{i\theta} \Leftrightarrow h\lambda = e^{i\theta} - 1, \quad \theta \in [0, 2\pi)$$



## Stability Regions for Higher Order Taylor Series Methods

Consider

$$y(t_{i+1}) = \sum_{k=0}^n \frac{h^k}{k!} y^{(k)}(t_i) + \frac{h^{n+1}}{(n+1)!} y^{(n+1)}(\xi_i), \quad \xi_i \in [t_i, t_{i+1}]$$

Now, with  $y'(t) = \lambda y(t)$  we have

$$y''(t) = \lambda y'(t) = \lambda^2 y(t)$$

So

$$y^{(n)}(t) = \lambda^n y(t)$$

And it follows that

$$y(t_{i+1}) = \sum_{k=0}^n \frac{(h\lambda)^k}{k!} y(t_i) + \frac{(h\lambda)^{n+1}}{(n+1)!} y^{(n+1)}(\xi_i), \quad \xi_i \in [t_i, t_{i+1}]$$

## Stability Regions for Higher Order Taylor Series Methods

The stability criterion is given by the relation

$$y(t_{i+1}) = y(t_0) \left[ \sum_{k=0}^n \frac{(h\lambda)^k}{k!} \right]^{i+1}$$

i.e.

$$\left| \sum_{k=0}^n \frac{(h\lambda)^k}{k!} \right| \leq 1$$

Again, the boundary is given by

$$\sum_{k=0}^n \frac{(h\lambda)^k}{k!} = e^{i\theta}$$


## Plotting the Boundary of the Stability Region

For  $n = 4$  we have

$$\frac{(h\lambda)^4}{24} + \frac{(h\lambda)^3}{6} + \frac{(h\lambda)^2}{2} + (h\lambda) + 1 - e^{i\theta} = 0$$

**matlab**)>> z=roots([1/24 1/6 1/2 1 1-exp(i\*theta)])

Now, vary  $\theta$  in the interval  $[0, 2\pi)$ , collect all the roots, and plot in the complex plane ( $x = \text{real}(z)$ ,  $y = \text{imag}(z)$ ) —



**Figure:** The circle corresponding to Euler's Method is included for comparison.


## Some Comments on Higher Order Taylor Series Methods

In the cases where the derivative(s)  $f^{(k)}(t, y)$  can be computed, higher order ( $n > 1$ ) Taylor series methods are superior to Euler's method (Taylor order 1) for two reasons:

- 1 The local truncation error is smaller  $\sim \mathcal{O}(h^n)$
- 2 The region of stability is larger(!), allowing for (slightly) larger step-sizes  $h$ .

Next slide shows the stability regions for Taylor's Method of orders 1 through 8.

## Regions of Stability for Taylor's Method ( $n = 1, 2, \dots, 8$ )



## Improving Euler's Method: Alternatives

When the derivative(s) of  $f(t, y)$  cannot be computed —  $f$  may be a result of measurements — and/or is too expensive to compute/evaluate, we need alternative approaches to improve on Euler's Method.

We are going to explore the following approaches:

- Runge-Kutta Methods
- Linear Multistep Methods
- Predictor-Corrector Methods

There is significant overlap between these different approaches, hence we will “re-discover” some methods in several contexts.

## Runge Kutta Methods

### Runge-Kutta (RK) methods

- One-step methods — moving from time  $t_n$  to time  $t_{n+1}$ : Still easy to build adaptive methods if/when necessary (step-length changes on-the-fly are “easy.”)
- Breaks/complicates linearity — the structure of the local error becomes more complicated.

**Catch-22:** Easy to change step-size since it is a one-step method, but hard(er) to tell when it is needed (local error complicated).

## Linear Multistep Methods: Reverse Catch-22

When we look at Linear Multistep Methods (which use multiple points  $y_n, y_{n-1}, \dots, y_{n-k}$  in order to compute  $y_{n+1}$ ), we will see that they have the reverse problem: —

- for this class of methods it is easy to estimate the local error ( $\Rightarrow$  easy to know when a change in step-size is necessary to maintain a certain level of local accuracy),
- but the multistep structure makes it hard to change the step-size...

## A General $s$ -stage RK method

A general  $s$ -stage RK method for the problem

$$y'(t) = f(t, y), \quad y(t_0) = y_0$$

is defined by

$$y_{n+1} = y_n + h \sum_{i=1}^s b_i k_i$$

where the  $k_i$ s are multiple estimates of the right-hand-side  $f(t, y)$

$$k_i = f \left( t_n + c_i h, y_n + h \sum_{j=1}^s a_{i,j} k_j \right), \quad i = 1, 2, \dots, s$$

with the following row-sum condition

$$c_i = \sum_{j=1}^s a_{i,j} \quad i = 1, 2, \dots, s$$

## Example: Heun's Method is an RK 2-stage Method

$$k_1 = f(t_n, y_n) \quad \Rightarrow c_1 = 0, a_{1,j} = 0$$

$$k_2 = f(t_n + h, y_n + h k_1) \quad \Rightarrow c_2 = 1, a_{2,1} = 1, a_{2,2} = 0$$

$$y_{n+1} = y_n + h \left[ \frac{k_1 + k_2}{2} \right] \quad \Rightarrow b_1 = b_2 = \frac{1}{2}$$

The **Butcher Array** describing Heun's Method

$$\begin{array}{c|cc} c_1 & a_{1,1} & a_{1,2} \\ c_2 & a_{2,1} & a_{2,2} \\ \hline & b_1 & b_2 \end{array} = \begin{array}{c|cc} 0 & 0 & 0 \\ 1 & 1 & 0 \\ \hline & 1/2 & 1/2 \end{array}$$

## Example: Euler's Midpoint Method

$$k_1 = f(t_n, y_n) \quad \Rightarrow c_1 = 0, a_{1,j} = 0$$

$$k_2 = f \left( t_n + \frac{h}{2}, y_n + \frac{k_1 h}{2} \right) \quad \Rightarrow c_2 = \frac{1}{2}, a_{2,1} = \frac{1}{2}, a_{2,2} = 0$$

$$y_{n+1} = y_n + h k_2 \quad \Rightarrow b_1 = 0, b_2 = 1$$

The **Butcher Array** describing Euler's Midpoint Method

$$\begin{array}{c|cc} c_1 & a_{1,1} & a_{1,2} \\ c_2 & a_{2,1} & a_{2,2} \\ \hline & b_1 & b_2 \end{array} = \begin{array}{c|cc} 0 & 0 & 0 \\ 1/2 & 1/2 & 0 \\ \hline & 0 & 1 \end{array}$$

## The Butcher Array

The Butcher array for a general  $s$ -stage RK method is

$$\begin{array}{c|cccc} c_1 & a_{1,1} & a_{1,2} & \cdots & a_{1,s} \\ c_2 & a_{2,1} & a_{2,2} & \cdots & a_{2,s} \\ \vdots & \vdots & & & \vdots \\ c_s & a_{s,1} & a_{s,2} & \cdots & a_{s,s} \\ \hline & b_1 & b_2 & \cdots & b_s \end{array} = \begin{array}{c|c} \tilde{\mathbf{c}} & A \\ \hline & \tilde{\mathbf{b}}^T \end{array}$$

We define the  $s$ -dimensional vectors  $\tilde{\mathbf{b}}$ ,  $\tilde{\mathbf{c}}$  and the  $s \times s$ -matrix  $A$ :

$$\tilde{\mathbf{b}} = [b_1, b_2, \dots, b_s]^T, \quad \tilde{\mathbf{c}} = [c_1, c_2, \dots, c_s]^T, \quad A = [a_{i,j}]_{i,j=1}^s$$

## 3 Types of RK-methods, I/III

- **Explicit** (e.g. Heun's and Midpoint):

- If each  $k_j$  only depends on previously computed  $k_i$  ( $i < j$ ), then the method is **explicit**, and the matrix  $A$  is **strictly lower triangular** (i.e. the elements on and above the diagonal are zero).

## 3 Types of RK-methods, II/III

- **Semi-implicit\***

- If  $A$  is lower-triangular with non-zero entries on the diagonal, then each  $k_i$  is defined by a non-linear system:

$$k_i = f \left( t_n + c_i h, y_n + \sum_{j=1}^i a_{i,j} k_j \right), \quad i = 1, 2, \dots, s$$

We have to solve  $s$  non-linear (but uncoupled) systems of equations in each iteration...

\* Butcher (1965) calls these methods "Semi-implicit," Norsett (1974) "Semi-explicit," and Alexander (1977) "Diagonally Implicit RK" or DIRK methods.

## 3 Types of RK-methods, III/III

- **Implicit:**

- If  $A$  is a general matrix (non-zeros above the diagonal) then each  $k_i$  is defined by a non-linear system:


$$k_i = f \left( t_n + c_i h, y_n + \sum_{j=1}^s a_{i,j} k_j \right), \quad i = 1, 2, \dots, s$$

We have to solve  $s$  non-linear **coupled** systems of equations in each iteration... This can be a daunting computational task (see *Math 693a*); most of the time we will try to avoid going this route!

RK-methods constitute a sensible idea. The unique solution to a well posed initial value ODE problem

$$y'(t) = f(t, y), \quad y(t_0) = y_0$$

is a single curve in  $(t, y)$ -space. Solutions to the same ODE with (slightly) different initial conditions form a family of solutions:



Due to numerical errors — truncation, and roundoff errors — any numerical solution “wanders off” the exact solution curve. The numerical solution is affected by neighboring solutions.

RK-methods gather information about this “family” of solution curves.

An explicit RK-method sends out “feelers” into solution space, gathering samples of the derivative, and then decides in what direction to take the final Euler-like step.

Paraphrased from J.D. Lambert, “*Numerical Solutions for Ordinary Differential Systems: the Initial Value Problem.*”

## Deriving Explicit 1-stage RK-methods

The Butcher array for an 1-stage RK method has the form:

$$\begin{array}{c|c} c_1 & a_{1,1} \\ \hline & b_1 \end{array}$$

If we want an explicit scheme, then  $a_{1,1} = 0$ , and since  $c_1 = \sum_{j=1}^1 a_{1,j}$ , we have  $c_1 = 0$ . Further **consistency\*** requires that  $\sum b_j = 1$ , so  $b_1 = 1$ . We are left with

$$\begin{array}{c|c} 0 & 0 \\ \hline & 1 \end{array}$$

or

$$\begin{aligned} k_1 &= f(t_n, y_n) \\ y_{n+1} &= y_n + hk_1 = y_n + hf(t_n, y_n), \quad \text{Euler's Method!} \end{aligned}$$

\* We have yet to prove this condition.

## Deriving Explicit 2-stage RK-methods, I/III

The Butcher array for a 2-stage explicit RK method has the form:

$$\begin{array}{c|cc} 0 & 0 & 0 \\ c_2 & a_{2,1} & 0 \\ \hline & b_1 & b_2 \end{array} \sim \begin{array}{c|cc} 0 & 0 & 0 \\ c_2 & c_2 & 0 \\ \hline & b_1 & 1 - b_1 \end{array}$$

Hence,

$$\begin{cases} k_1 = f(t_n, y_n) \\ k_2 = f(t_n + c_2 h, y_n + c_2 h k_1) \\ y_{n+1} = y_n + h [b_1 k_1 + (1 - b_1) k_2] \end{cases}$$

describes all possible explicit 2-stage RK-methods.

How do we choose the parameters  $c_2$  and  $b_1$ ???

— Taylor Expansion, of course!

## Deriving Explicit 2-stage RK-methods, II/III

With the following Taylor expansions:

$$\begin{aligned} y_{n+1} &= y_n + hf_n + \frac{h^2}{2} f'_n + \mathcal{O}(h^3) \\ k_1 &= f_n \\ k_2 &= f(t_n + c_2h, y_n + c_2hk_1) \\ &= f_n + (c_2h) \frac{\partial}{\partial t} f(t_n, y_n) + (c_2hf_n) \frac{\partial}{\partial y} f(t_n, y_n) + \mathcal{O}(h^2) \end{aligned}$$

We can define the Local Truncation Error

$$\begin{aligned} \text{LTE}(h) &= \frac{y_{n+1} - y_n}{h} - b_1k_1 - (1 - b_1)k_2 \\ &= \left[ f_n + \frac{h}{2} f'_n + \mathcal{O}(h^2) \right] - \\ &\quad - \left[ b_1f_n - (1 - b_1) \left( f_n + (c_2h) \left[ \frac{\partial}{\partial t} f_n + \frac{\partial}{\partial y} f_n \cdot f_n \right] \right) \right] \\ &= \frac{h}{2} \left[ \frac{\partial}{\partial t} f_n + \frac{\partial}{\partial y} f_n \cdot f_n \right] - \mathbf{b}_2c_2h \left[ \frac{\partial}{\partial t} f_n + \frac{\partial}{\partial y} f_n \cdot f_n \right] + \mathcal{O}(h^2) \end{aligned}$$

## Deriving Explicit 2-stage RK-methods, III/III

We have

$$\text{LTE}(h) = \frac{h}{2} \left[ \frac{\partial}{\partial t} f_n + \frac{\partial}{\partial y} f_n \cdot f_n \right] - \mathbf{b}_2c_2h \left[ \frac{\partial}{\partial t} f_n + \frac{\partial}{\partial y} f_n \cdot f_n \right] + \mathcal{O}(h^2)$$

Now, if

$$\frac{h}{2} - b_2c_2h = 0 \quad \Leftrightarrow 2b_2c_2 = 1$$

we get  $\text{LTE}(h) \sim \mathcal{O}(h^2)$ , i.e. our 2-stage RK-method is **second order**.

The corresponding family of Butcher arrays is

$$\begin{array}{c|cc} 0 & 0 & 0 \\ c_2 & c_2 & 0 \\ \hline & 1 - 1/(2c_2) & 1/(2c_2) \end{array}$$

Sanity check:  $c_2 = 1/2$  gives Euler's Midpoint Method, and  $c_2 = 1$  gives Heun's Method.

## Deriving Explicit Higher Order RK-methods

We can use the same approach — Taylor expansion and parameter matching, to find higher order explicit RK-methods.

**Natural question:** Is this the best way of deriving the RK-methods?

**Answer:** There are more elegant methods for deriving the RK-methods. Most of the work was done by Butcher starting in the mid-1960s. The methods depend on defining the **Frechet derivative** and also requires some basic understanding of graph theory ("rooted trees.")

Butcher, J.C. (1987), *The Numerical Analysis of Ordinary Differential Equations: Runge-Kutta and General Linear Methods*, Wiley, Chichester.

## Example: 3-stage RK-method

$$\begin{array}{c|ccc} c_1 & a_{1,1} & a_{1,2} & a_{1,3} \\ c_2 & a_{2,1} & a_{2,2} & a_{2,3} \\ c_3 & a_{3,1} & a_{3,2} & a_{3,3} \\ \hline & b_1 & b_2 & b_3 \end{array} = \begin{array}{c|ccc} 0 & 0 & 0 & 0 \\ 1/2 & 1/2 & 0 & 0 \\ 1 & -1 & 2 & 0 \\ \hline & 1/6 & 2/3 & 1/6 \end{array}$$

$$k_1 = f(t_n, y_n)$$

$$k_2 = f\left(t_n + \frac{h}{2}, y_n + \frac{hk_1}{2}\right)$$


$$k_3 = f(t_n + h, y_n - hk_1 + 2hk_2)$$

$$y_{n+1} = y_n + \frac{h}{6} (k_1 + 4k_2 + k_3)$$

(See next slide for visualization.)

### Example: 3-stage RK-method

### One Step



**Figure:** Different "stages" of one step of the 3-stage RK method.

### Example: 4-stage RK-method (Attributed to Runge)

$$\begin{array}{c|cccc|cccc}
 c_1 & a_{1,1} & a_{1,2} & a_{1,3} & a_{1,4} & 0 & 0 & 0 & 0 \\
 c_2 & a_{2,1} & a_{2,2} & a_{2,3} & a_{2,4} & 1/2 & 1/2 & 0 & 0 \\
 c_3 & a_{3,1} & a_{3,2} & a_{3,3} & a_{3,4} & 1/2 & 0 & 1/2 & 0 \\
 c_4 & a_{4,1} & a_{4,2} & a_{4,3} & a_{4,4} & 1 & 0 & 0 & 1 \\
 \hline
 & b_1 & b_2 & b_3 & b_4 & & 1/6 & 1/3 & 1/3 & 1/6
 \end{array} =$$

$$k_1 = f(t_n, y_n)$$

$$k_2 = f\left(t_n + \frac{h}{2}, y_n + \frac{hk_1}{2}\right)$$


$$k_3 = f\left(t_n + \frac{h}{2}, y_n + \frac{hk_2}{2}\right)$$

$$k_4 = f(t_n + h, y_n + hk_3)$$

$$y_{n+1} = y_n + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4)$$

### Example: Runge's 4-stage method

### One Step



### Example: 4-stage RK-method (Attributed to Kutta)

$$\begin{array}{c|cccc|cccc}
 c_1 & a_{1,1} & a_{1,2} & a_{1,3} & a_{1,4} & 0 & 0 & 0 & 0 \\
 c_2 & a_{2,1} & a_{2,2} & a_{2,3} & a_{2,4} & 1/3 & 1/3 & 0 & 0 \\
 c_3 & a_{3,1} & a_{3,2} & a_{3,3} & a_{3,4} & 2/3 & -1/3 & 1 & 0 \\
 c_4 & a_{4,1} & a_{4,2} & a_{4,3} & a_{4,4} & 1 & 1 & -1 & 1 \\
 \hline
 & b_1 & b_2 & b_3 & b_4 & & 1/8 & 3/8 & 3/8 & 1/8
 \end{array} =$$

$$k_1 = f(t_n, y_n)$$


$$k_2 = f\left(t_n + \frac{h}{3}, y_n + \frac{hk_1}{3}\right)$$

$$k_3 = f\left(t_n + \frac{2h}{3}, y_n - \frac{hk_1}{3} + hk_2\right)$$

$$k_4 = f(t_n + h, y_n + hk_1 - hk_2 + hk_3)$$

$$y_{n+1} = y_n + \frac{h}{8}(k_1 + 3k_2 + 3k_3 + k_4)$$





Stability Regions Revisited & RK-Methods — (33/34)

We have some important residual issues related to Runge-Kutta methods to clear up next time:

- **Consistency** condition: The requirement  $\sum b_i = 1$ .
- **Error estimation** (using Richardson's Extrapolation).
- **Stability Analysis**.
- Some more examples of RK-methods in action.

Future topic:

- Deriving RK-methods using rooted trees.