

Numerical Solutions to Differential Equations

Lecture Notes #11 — Runge-Kutta Methods for Stiff ODEs

Peter Blomgren,
(blomgren.peter@gmail.com)

Department of Mathematics and Statistics
Dynamical Systems Group
Computational Sciences Research Center
San Diego State University
San Diego, CA 92182-7720
<http://terminus.sdsu.edu/>

Spring 2015

Outline

1 Introduction

- Review: Stability for Explicit Runge-Kutta Methods
- Stability of Semi-Implicit RK-Methods

2 Approximations of e^x

- Optimal Polynomial Approximations
- Optimal Rational (Padè) Approximations
- Rational Approximations: Classification and Properties

3 Implicit RK-Methods for Stiff Problems

- Examples: Gauss-Legendre Methods
- Wishing for L-stability... The Radau Methods

Recall: Stability Analysis for Explicit RK-methods

By applying the RK-methods to the scalar test-problem $\mathbf{y}'(\mathbf{t}) = \lambda \mathbf{y}(\mathbf{t})$, $\mathbf{y}(\mathbf{t}_0) = \mathbf{y}_0$ we will find the regions of stability for the methods.

E.g. Heun's Method

$$\begin{array}{c|cc} c_1 & a_{1,1} & a_{1,2} \\ c_2 & a_{2,1} & a_{2,2} \\ \hline & b_1 & b_2 \end{array} = \begin{array}{c|cc} 0 & 0 & 0 \\ 1 & 1 & 0 \\ \hline & 1/2 & 1/2 \end{array}$$

Hence

$$k_1 = f(t_n, y_n) = \lambda y_n$$

$$k_2 = f(t_n + h, y_n + h k_1) = \lambda(y_n + h k_1) = \lambda y_n + h \lambda^2 y_n$$

$$y_{n+1} = y_n \left[1 + \frac{h}{2} [2\lambda + h\lambda^2] \right] = y_n \underbrace{\left[1 + h\lambda + \frac{(h\lambda)^2}{2} \right]}_{R(h\lambda)}$$

Recall: Stability of Heun's Method

1/11

The iteration is given by

$$y_{n+1} = R(h\lambda)y_n,$$

and the stability region is given by

$$|R(h\lambda)| = \left| 1 + h\lambda + \frac{(h\lambda)^2}{2} \right| \leq 1.$$

We find the boundary of the region by find the complex roots of

$$1 - e^{i\theta} + h\lambda + \frac{(h\lambda)^2}{2} = 0,$$


for all values of $\theta \in [0, 2\pi)$.

Recall: Stability of Heun's Method

11/11

We find the boundary of the region by find the complex roots of

$$1 - e^{i\theta} + h\lambda + \frac{(h\lambda)^2}{2} = 0, \quad \forall \theta \in [0, 2\pi).$$



Recall: Stability Regions for General RK-methods

1/11

For notational convenience we absorb $h\lambda \rightarrow \hat{h}$.

Using the A from the Butcher array, we can write the k_i 's

$$\tilde{\mathbf{k}} = \begin{bmatrix} k_1 \\ k_2 \\ \vdots \\ k_s \end{bmatrix} = y_n \tilde{\mathbf{1}} + \hat{h} A \tilde{\mathbf{k}}, \quad \text{where } \tilde{\mathbf{1}} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix},$$

thus, we can solve for $\tilde{\mathbf{k}}$:

$$\tilde{\mathbf{k}} = (I - \hat{h}A)^{-1} \tilde{\mathbf{1}} y_n.$$

Further,

$$y_{n+1} = y_n + \hat{h} \tilde{\mathbf{b}}^T \tilde{\mathbf{k}} = y_n + \hat{h} \tilde{\mathbf{b}}^T (I - \hat{h}A)^{-1} \tilde{\mathbf{1}} y_n.$$

Recall: Stability Regions for General RK-methods

11/11

We have

$$y_{n+1} = y_n + \hat{h}\tilde{\mathbf{b}}^T \tilde{\mathbf{k}} = y_n + \hat{h}\tilde{\mathbf{b}}^T (I - \hat{h}A)^{-1} \tilde{\mathbf{l}} y_n.$$

Thus, the stability function is

Stability Function, $R(\hat{h})$

$$R(\hat{h}) = 1 + \hat{h}\tilde{\mathbf{b}}^T (I - \hat{h}A)^{-1} \tilde{\mathbf{l}}.$$

As usual, the method is stable for \hat{h} such that $|R(\hat{h})| \leq 1$.

For explicit methods, A strictly lower triangular, the quantity

$$\tilde{\mathbf{d}} = (I - \hat{h}A)^{-1} \tilde{\mathbf{l}},$$

is easily computable using forward substitution.

The Stability Function $R(\hat{h})$

As we have seen, the stability functions for explicit RK-methods are **polynomials**... Lets consider the stability analysis for a **semi-implicit** method defined by the following Butcher array:

$$\begin{array}{c|cc} c_1 & a_{1,1} & 0 \\ c_2 & a_{2,1} & a_{2,2} \\ \hline & b_1 & b_2 \end{array}$$

We get

$$\begin{aligned} k_1 &= f(t_n + c_1 h, y_n + h k_1 a_{1,1}) &= \lambda y_n + \hat{h} a_{1,1} k_1 \\ k_2 &= f(t_n + c_2 h, y_n + h k_1 a_{2,1} + h k_2 a_{2,2}) &= \lambda y_n + \hat{h} (a_{2,1} k_1 + a_{2,2} k_2) \end{aligned}$$

$$k_1 = \left[\frac{1}{1 - \hat{h} a_{1,1}} \right] \lambda y_n, \quad k_2 = \left[\frac{1 - \hat{h} a_{1,1} - \hat{h} a_{2,1}}{(1 - \hat{h} a_{1,1})(1 - \hat{h} a_{2,2})} \right] \lambda y_n.$$

The Stability Function $R(\hat{h})$ — Semi Implicit RK

With these values of k_1, k_2 :

$$k_1 = \left[\frac{1}{1 - \hat{h}a_{1,1}} \right] \lambda y_n, \quad k_2 = \left[\frac{1 - \hat{h}a_{1,1} - \hat{h}a_{2,1}}{(1 - \hat{h}a_{1,1})(1 - \hat{h}a_{2,2})} \right] \lambda y_n$$

the final step becomes

$$\begin{aligned} y_{n+1} &= y_n [1 + hb_1k_1 + hb_2k_2] \\ &= y_n \underbrace{\left[1 + \hat{h} \left[\frac{b_1}{1 - \hat{h}a_{1,1}} + \frac{b_2(1 - \hat{h}a_{1,1} - \hat{h}a_{2,1})}{(1 - \hat{h}a_{1,1})(1 - \hat{h}a_{2,2})} \right] \right]}_{R(\hat{h})} \end{aligned}$$

Clearly, $R(\hat{h})$ is a rational function.

Summarizing...

1/11

We have seen that when we apply an RK-method to the test equation $y'(t) = \lambda y(t)$, we get the discrete iteration

$$y_{n+1} = R(\hat{h})y_n, \quad \hat{h} = h\lambda, \quad \lambda \in \mathbb{C},$$

where

- for explicit RK-methods $R(\hat{h})$ is a polynomial, and
- for semi- (and fully) implicit RK-methods it is a rational function.

Summarizing...

11/11

The exact solution to the test equation is

$$\mathbf{y}(\mathbf{t}) = \mathbf{K}e^{\lambda \mathbf{t}}, \quad K \text{ constant (initial conditions)}$$

hence, the exact solution to the iteration is

$$y_{n+1}^* = e^{\lambda h} y_n = e^{\hat{h}} y_n.$$

We can express the truncation error as:

$$\text{LTE}(\hat{h}) = \frac{y_{n+1}^* - y_{n+1}}{h} = \frac{1}{h} \left[e^{\hat{h}} - R(\hat{h}) \right] y_n = \mathcal{O}(\hat{h}^p),$$

for a p^{th} order method.

Polynomial Approximations to the Exponential

Clearly the truncation error

$$\text{LTE}(\hat{h}) = \frac{1}{h} \left[e^{\hat{h}} - R(\hat{h}) \right] y_n = \mathcal{O}(\hat{h}^p)$$



only depends on how well $R(\hat{h})$ approximates the exponential $e^{\hat{h}}$!!!

Hence, if we know how to find a good approximation to the exponential, we can back-track and build a high-order scheme (hopefully with good stability properties).

The optimal polynomial approximations come directly from the **Taylor expansion** of $e^{\hat{h}}$:

$$e^{\hat{h}} = \sum_{k=0}^{\infty} \frac{1}{k!} \hat{h}^k.$$

Rational Approximations to the Exponential

I/II

We are now motivated to look at **Rational Approximations to the Exponential**^{Math 541}.

Value-Add (Strong Connection to Stability)

The value-add is that we are working directly with the stability function. Once we find high-order approximations to $e^{\hat{h}}$ with desirable stability properties we go back and identify coefficients to build the corresponding finite-difference scheme.

Let

$$R_T^S(\hat{h}) = \left[\sum_{i=0}^S a_i \hat{h}^i \right] / \left[\sum_{j=0}^T b_j \hat{h}^j \right] \quad a_0 = b_0 = 1, a_S \neq 0, b_T \neq 0$$

denote a rational approximation of $e^{\hat{h}}$.

Rational Approximations to the Exponential

II/II

The maximum order of approximation of the exponential for a rational function $R_T^S(\hat{h})$ is $T + S$:

$$e^{\hat{h}} - R_T^S(\hat{h}) = \mathcal{O}(\hat{h}^{p+1}), \quad p \leq T + S$$

if $p = S + T$ then $R_T^S(\hat{h})$ is called a **Padé Approximation** of $e^{\hat{h}}$.

Butcher (1987) figured out what the coefficients for the Padé approximations (of e^x) are:

$$a_i = \frac{S!}{(S+T)!} \frac{(S+T-i)!}{i!(S-i)!}, \quad i = 1, 2, \dots, S$$

$$b_j = (-1)^j \frac{T!}{(S+T)!} \frac{(S+T-j)!}{j!(T-j)!}, \quad j = 1, 2, \dots, T$$

Examples: Some Padé Approximations — Order 3

$$R_3^0(\hat{h}) = \frac{1}{1 - \hat{h} + \frac{1}{2}\hat{h}^2 - \frac{1}{6}\hat{h}^3}$$

$$R_2^1(\hat{h}) = \frac{1 + \frac{1}{3}\hat{h}}{1 - \frac{2}{3}\hat{h} + \frac{1}{6}\hat{h}^2}$$


$$R_1^2(\hat{h}) = \frac{1 + \frac{2}{3}\hat{h} + \frac{1}{6}\hat{h}^2}{1 - \frac{1}{3}\hat{h}}$$

$$R_0^3(\hat{h}) = 1 + \hat{h} + \frac{1}{2}\hat{h}^2 + \frac{1}{6}\hat{h}^3$$


As usual, the boundaries of the stability regions are given by

$$R_T^S(\hat{h}) = e^{i\theta}, \quad \theta \in [0, 2\pi)$$


The Associated Stability Regions




$R^3_0(\hat{h})$ — interior



$R^2_1(\hat{h})$ — interior







$R^1_2(\hat{h})$ — exterior



$R^0_3(\hat{h})$ — exterior


The Associated Stability Regions with Magnitude

Order 3









The Associated Stability Regions with Magnitude

Order 4



The Associated Stability Regions with Magnitude

Order 5



Definition: Acceptability of Approximation

Definition (Ehle, 1969)

A rational approximation $R(\hat{h})$ to $e^{\hat{h}}$ is said to be:

- ❶ **A-acceptable** if $|R(\hat{h})| < 1$ whenever $\operatorname{Re}(\hat{h}) < 0$.
- ❷ **A_0 -acceptable** if $|R(\hat{h})| < 1$ whenever \hat{h} is real and negative.
- ❸ **L-acceptable** if it is A-acceptable, and $|R(\hat{h})| \rightarrow 0$ as $\operatorname{Re}(\hat{h}) \rightarrow -\infty$.

Clearly the associated numerical methods are A-stable, A_0 -stable, and L -stable.

Theorems: Acceptability of Padé Approximations

Theorem (Varga, 1961)

If $T \geq S$, then $R_T^S(\hat{h})$ is A_0 -acceptable.

Theorem (Birkhoff and Varga, 1965)

If $T = S$, then $R_T^S(\hat{h})$ is A -acceptable.

Theorem (Ehle, 1969)


If $T = S + 1$, or $T = S + 2$ then $R_T^S(\hat{h})$ is L -acceptable.

Theorem (Wanner, Hairer, Nørsett, 1978)


*$R_T^S(\hat{h})$ is A -acceptable if and only if $T - 2 \leq S \leq T$.
("The Ehle Conjecture" 1965)*

Theorems — Visualized

Guaranteed A_0 -acceptability of $R_T^S(h)$, Varga 1961




Guaranteed A-acceptability of $R_T^S(h)$, Birkhoff and Varga 1965




Theorems — Visualized


Guaranteed L-acceptability of $R_T^S(h)$, Ehle 1969



A-acceptability of $R_T^S(h)$, W-H-N 1978



Theorems — Note



Note: Even though $\lim_{\text{Real}(\hat{h}) \rightarrow -\infty} |R_3^0(\hat{h})| \rightarrow 0$, $R_3^0(\hat{h})$ is not L-acceptable, since it is **not** A-acceptable; — The left-half-plane of the region of absolute stability has two small “cut-outs.” It is $A(\alpha)$ -acceptable, where $\alpha \approx \frac{\pi}{2} - 0.031$.

Implicit RK-methods Suitable for Stiff Systems

Given the preceding detour into approximation of the exponential, we are now ready to take another look at RK-methods.



Given an RK-method, with its associated Butcher array

$$\begin{array}{c|c} \tilde{\mathbf{c}} & A \\ \hline & \tilde{\mathbf{b}}^T \end{array}$$

we recall that we can express the stability function as

$$R(\hat{h}) = 1 + \hat{h}\tilde{\mathbf{b}}^T(I - \hat{h}A)^{-1}\tilde{\mathbf{1}},$$

or

$$R(\hat{h}) = \frac{\det[I - \hat{h}(A - \tilde{\mathbf{1}}\tilde{\mathbf{b}}^T)]}{\det[I - \hat{h}A]}.$$

Finding the RK-method from $R(\hat{h})$

- Whereas it is possible, in some cases (but extremely tedious, in all cases) to take a rational function $R(\hat{h})$ and “reverse engineer” a numerical method, this is not the path we will take.
- We are going to look at the fully implicit **Gauss** or **Gauss-Legendre Methods**:
- By **optimally** selecting the points where f is evaluated (the entries in the matrix A which occurs in the Butcher array), an s -stage Gauss method achieves order $2s$.

Note: The optimal placement of the (time, \vec{c}) points comes directly from the analysis for Gaussian numerical integration ^{Math 541}.

Gauss(-Legendre) Methods

I/III

Since there is a **unique** $R_S^S(\hat{h})$ rational approximation to order $2s$ of $e^{\hat{h}}$, namely the Padé approximation, it follows that the stability function for the Gauss methods must be the Padé approximation.

Since $S = T$ all Gauss methods are A-stable (Birkhoff-Varga).

Example (“Implicit Mid-point Rule.”)

The “Implicit Mid-point Rule” is a 1-stage 2nd-order Gauss method:

$$\begin{array}{c|c} \frac{1}{2} & \frac{1}{2} \\ \hline & 1 \end{array}$$

$$y_{n+1} = y_n + hf \left(t_n + \frac{1}{2}h, \frac{1}{2}(y_n + y_{n+1}) \right)$$

Gauss(-Legendre) Methods

II/III

Example (2-stage 4th order Gauss method)

$\frac{3-\sqrt{3}}{6}$	$\frac{1}{4}$	$\frac{3-2\sqrt{3}}{12}$
$\frac{3+\sqrt{3}}{6}$	$\frac{3+2\sqrt{3}}{12}$	$\frac{1}{4}$
	$\frac{1}{2}$	$\frac{1}{2}$

Gauss(-Legendre) Methods



Example (3-stage 6th order Gauss method)





$\frac{5-\sqrt{15}}{10}$	$\frac{5}{36}$	$\frac{10-3\sqrt{15}}{45}$	$\frac{25-6\sqrt{15}}{180}$
$\frac{1}{2}$	$\frac{10+3\sqrt{15}}{72}$	$\frac{2}{9}$	$\frac{10-3\sqrt{15}}{72}$
$\frac{5+\sqrt{15}}{10}$	$\frac{25+6\sqrt{15}}{180}$	$\frac{10+3\sqrt{15}}{45}$	$\frac{5}{36}$
	$\frac{5}{18}$	$\frac{4}{9}$	$\frac{5}{18}$

Ponder how much fun would it be to reverse engineer this 3-6 method from the Padé approximation

$$R_3^3(\hat{h}) = \frac{1 + \frac{1}{2}\hat{h} + \frac{1}{10}\hat{h}^2 + \frac{1}{120}\hat{h}^3}{1 - \frac{1}{2}\hat{h} + \frac{1}{10}\hat{h}^2 - \frac{1}{120}\hat{h}^3}$$

Gauss(-Legendre) Methods

Stability



Gauss(-Legendre) Methods — The Final Wish

- If want to find something “wrong” with the Gauss methods, it would be that they are **not L-stable**.
- It turns out we can trade one order of approximation for L-stability. The **Radau I-A** and **Radau II-A** s -stage methods are order $(2s - 1)$ and L-stable.
- The Radau I-A methods are derived just like the Gaussian methods, but require the left endpoint to be part of the interval ($c_1 = 0$).
- The Radau II-A methods require the right endpoint to be part of the interval ($c_s = 1$).

Radau I/II-A Methods

Examples I/II

Example (1-stage 1st order Radau II-A L-stable method)

$$\begin{array}{c|c} 1 & 1 \\ \hline & 1 \end{array}$$

Example (2-stage 3rd order Radau I-A L-stable method)

$$\begin{array}{c|cc} 0 & \frac{1}{4} & -\frac{1}{4} \\ \frac{2}{3} & \frac{1}{4} & \frac{5}{12} \\ \hline & \frac{1}{4} & \frac{3}{4} \end{array}$$




Radau I/II-A Methods

Examples II/II

Example (3-stage 5th order Radau I-A L-stable method)

0	$\frac{1}{9}$	$\frac{-1-\sqrt{6}}{18}$	$\frac{-1+\sqrt{6}}{18}$
$\frac{6-\sqrt{6}}{10}$	$\frac{1}{9}$	$\frac{88+7\sqrt{6}}{360}$	$\frac{88-43\sqrt{6}}{360}$
$\frac{6+\sqrt{6}}{10}$	$\frac{1}{9}$	$\frac{88+43\sqrt{6}}{360}$	$\frac{88-7\sqrt{6}}{360}$
	$\frac{1}{9}$	$\frac{16+\sqrt{6}}{36}$	$\frac{16-\sqrt{6}}{36}$

Radau Methods — Some Stability Regions Visualized



RK-methods — Wrap-up

I/II

- Clearly, constructing A- or L-stable implicit RK-methods is not an insurmountable task.
- Further, implementing the methods is also quite straight-forward.
- Either with the help of Richardson Extrapolation or by RKF45-like methods we can get good error estimates, and thus construct adaptive algorithms that change the step-size h on the fly.

RK-methods — Wrap-up

11/11

- These methods will work and can be designed to be very robust.
- However, in terms of **efficiency** they fall short of fine-tuned BDF (LMM) methods.
- To make RK-methods competitive, the computational handling of the implicitness must be cut down. There are a number of “tricks” — transformations that can be applied to reduce the computational burden.

Next couple of lectures...

- Linear Multistep Methods for Stiff ODEs.
- Review and examples.
- Hybrid Methods.
- Tie up loose ends.
- Start thinking about projects....