

Numerical Matrix Analysis

Notes #5 — The Singular Value Decomposition

Peter Blomgren
(blomgren@sdsu.edu)

Department of Mathematics and Statistics
Dynamical Systems Group
Computational Sciences Research Center
San Diego State University
San Diego, CA 92182-7720

<http://terminus.sdsu.edu/>

Spring 2024

(Revised: February 1, 2024)



Outline

- 1 Student Learning Targets, and Objectives
 - SLOs: The Singular Value Decomposition
- 2 Recap
 - Vector and Matrix Norm Inequalities
 - Missing Proof
- 3 Existence and Uniqueness of the SVD
 - The Theorem
 - Proof
- 4 The SVD
 - “Every Matrix is Diagonal”
 - Singular Values and Eigenvalues
 - The SVD \rightsquigarrow Matrix Properties

Student Learning Targets, and Objectives

Target The Singular Value Decomposition

Objective Existence and Uniqueness statements

Objective Impact: “diagonalizability”

Target The SVD \leftrightarrow Matrix Properties


Objective rank, range, null-space, norms

Objective relation to eigenvalues, determinant

Objective Linearly Optimal **Low Rank Approximations**

Gratuitous "AI"

Google Bard, 2004-02-01: create an image of a nerdy mathematician preparing slides for a lecture on computational matrix algebra



"Sup, bruh?!? I invert giant matrices by hand. What's your superpower?!?"

Last Time

- Hölder and Cauchy-[Bunyakovsky]-Schwarz inequalities:

$$|\vec{v}^* \vec{w}| \stackrel{H}{\leq} \|\vec{v}\|_p \|\vec{w}\|_q, \quad \frac{1}{p} + \frac{1}{q} = 1, \quad |\vec{v}^* \vec{w}| \stackrel{CBS}{\leq} \|\vec{v}\|_2 \|\vec{w}\|_2$$

- Bounds on the norms of matrix products

$$\|AB\| \leq \|A\| \|B\|$$

- General matrix norms: The Frobenius norm $\|A\|_F^2 = \sum_{ij} |a_{ij}|^2$.
- A geometrical introduction to the SVD.
- The reduced vs. the full SVD.

Missing Proof

We ended last lecture with: **“If** we can show that every matrix A has a SVD, **then** it follows that the image of the unit sphere under any linear map is a hyper-ellipse...”

We now turn our attention to showing that this indeed is the case...




Figure: The unit-sphere S^2 , and the image AS^2 , where $A = \begin{bmatrix} 1.3127 & 0.6831 & 0.6124 \\ 0.0129 & 1.0928 & 0.6085 \\ 0.3840 & 0.0353 & 1.0158 \end{bmatrix}$.

Theorem: $A = U\Sigma V^*$

Existence and Uniqueness

Theorem (Existence and Uniqueness of the SVD)

Every matrix $A \in \mathbb{C}^{m \times n}$ has a singular value decomposition $A = U\Sigma V^*$, where

$U \in \mathbb{C}^{m \times m}$ is unitary

$V \in \mathbb{C}^{n \times n}$ is unitary

$\Sigma \in \mathbb{R}^{m \times n}$ is diagonal, non-negative.

Furthermore, the singular values $\{\sigma_k\}$ are uniquely determined, and if A is square and the σ_k are *distinct*, the left $\{\vec{u}_k\}$ and right $\{\vec{v}_k\}$ singular vectors are uniquely determined up to complex scalar factors $s \in \mathbb{C} : |s| = 1$.

We present a proof that is very “matrix-y,” a completely different approach is presented [MATH 524 (NOTES#7.1–7.2)]

Theorem: $A = U\Sigma V^*$

Proof, 1 of 5

THE PROOF IS BY INDUCTION. Let $\sigma_1 = \|A\|_2$. There must exist $\vec{u}_1 \in \mathbb{C}^m$, $\|\vec{u}_1\|_2 = 1$, and $\vec{v}_1 \in \mathbb{C}^n$, $\|\vec{v}_1\|_2 = 1$, such that $A\vec{v}_1 = \sigma_1\vec{u}_1$:

$$\sigma_1 = \frac{\|A\vec{x}^*\|_2}{\|\vec{x}^*\|_2}, \text{ for some } \vec{x}^*. \quad \text{Let } \vec{v}_1 = \frac{\vec{x}^*}{\|\vec{x}^*\|_2}.$$

Clearly, $A\vec{v}_1 = \vec{p}$, for some \vec{p} . Let $\vec{u}_1 = \frac{\vec{p}}{\|\vec{p}\|_2}$, and $\sigma_1 = \|\vec{p}\|_2$.

Consider any extension (\exists Movie, see also [MATH 524]) of \vec{v}_1 to an orthonormal basis $\{\vec{v}_k\}_{k=1,\dots,n}$ of \mathbb{C}^n and of \vec{u}_1 to an orthonormal basis $\{\vec{u}_k\}_{k=1,\dots,m}$ of \mathbb{C}^m . Let U_1 and V_1 denote the matrices with columns \vec{u}_k and \vec{v}_k , respectively.

Theorem: $A = U\Sigma V^*$

Proof, 2 of 5

We have (by construction)

$$U_1^* A V_1 = S = \begin{bmatrix} \sigma_1 & \vec{w}^* \\ \vec{0} & B \end{bmatrix},$$

where $\vec{0}$ is a column-vector of size $(m-1)$, and \vec{w}^* is a row vector of size $(n-1)$, and the matrix $B \in \mathbb{C}^{(m-1) \times (n-1)}$.

Now,

$$\left\| \begin{bmatrix} \sigma_1 & \vec{w}^* \\ \vec{0} & B \end{bmatrix} \begin{bmatrix} \sigma_1 \\ \vec{w} \end{bmatrix} \right\|_2 \geq \sigma_1^2 + \vec{w}^* \vec{w} = \sqrt{\sigma_1^2 + \vec{w}^* \vec{w}} \left\| \begin{bmatrix} \sigma_1 \\ \vec{w} \end{bmatrix} \right\|_2,$$

Hence, $\|S\|_2 \geq \sqrt{\sigma_1^2 + \vec{w}^* \vec{w}}$.

Theorem: $A = U\Sigma V^*$

Proof, 3 of 5

We have $\|S\|_2 \geq \sqrt{\sigma_1^2 + \vec{w}^* \vec{w}}$, and $S = U_1^* A V_1$. Since U_1 and V_1 are unitary, we must have $\|S\|_2 = \|A\|_2 = \sigma_1$.

Therefore $\|\vec{w}\|_2^2 = \vec{w}^* \vec{w} = 0$, which means $\vec{w} = 0$, hence

$$U_1^* A V_1 = S = \begin{bmatrix} \sigma_1 & \vec{0}^* \\ \vec{0} & B \end{bmatrix}, \quad \Leftrightarrow \quad A = U_1 \begin{bmatrix} \sigma_1 & \vec{0}^* \\ \vec{0} & B \end{bmatrix} V_1^*$$

If $m = 1$, or $n = 1$, we are done. Otherwise, the sub-matrix B describes the action of A on the subspace orthogonal to \vec{v}_1 .

We can now recursively (inductively) apply the same process to B , and establish existence of the SVD of A :

$$A = U_1 \begin{bmatrix} 1 & \vec{0}^* \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} \sigma_1 & \vec{0}^* \\ \vec{0} & \Sigma_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0}^* \\ \vec{0} & V_2 \end{bmatrix}^* V_1^* = U \Sigma V^*.$$

Theorem: $A = U\Sigma V^*$

Proof, 4 of 5

THE UNIQUENESS PROOF REMAINS —

[Geometric Version] If the singular values σ_k are distinct, then the lengths of the semi-axes of the hyper-ellipse $A\mathbb{S}^{(n-1)}$ must be distinct.

The semi-axes themselves are determined by the geometry, up to a complex sign. $\square_{\text{geometric}}$.

[Algebraic Version] $\sigma_1 = \|A\|_2$ is uniquely determined. Now, suppose that in addition to \vec{v}_1 , there is another linearly independent vector \vec{w}_1 with $\|\vec{w}_1\| = 1$, and $\|A\vec{w}_1\| = \sigma_1$.

We define a unit vector \vec{v}_2 , orthogonal to \vec{v}_1 , as a linear combination of \vec{v}_1 and \vec{w}_1 :

$$\vec{v}_2 = \frac{\vec{w}_1 - (\vec{v}_1^* \vec{w}_1) \vec{v}_1}{\|\vec{w}_1 - (\vec{v}_1^* \vec{w}_1) \vec{v}_1\|_2}. \quad (\vec{v}_2 = \vec{w}_1 \perp \vec{v}_1)$$

Theorem: $A = U\Sigma V^*$

Proof, 5 of 5

Since $\|A\|_2 = \sigma_1$, $\|A\vec{v}_2\|_2 \leq \sigma_1$; but this must be an equality, otherwise since for some θ

$$\vec{w}_1 = \cos(\theta)\vec{v}_1 + \sin(\theta)\vec{v}_2, \quad \vec{v}_1 \perp \vec{v}_2, \quad \cos^2(\theta) + \sin^2(\theta) = 1$$

we would have $\|A\vec{w}_1\|_2 < \sigma_1$.

This vector \vec{v}_2 is a **second** right singular vector corresponding to the singular value σ_1 ; it will lead to the appearance of a \vec{y} (the last $(n-1)$ elements of $V_1^*\vec{v}_2$) with $\|\vec{y}\|_2 = 1$, and $\|B\vec{y}\|_2 = \sigma_1$.

Hence, if the singular vector \vec{v}_1 is not unique, then the corresponding singular value σ_1 is not simple ($\sigma_1 \neq \sigma_2$). Therefore there cannot exist a vector \vec{w}_1 as above.

Now, the uniqueness of the remaining singular vectors follows by induction. $\square_{\text{algebraic}}$

The SVD: $A = U\Sigma V^*$

Bold Statement

SVD enables us to say that **every matrix is "diagonal"** — as long as we use the proper bases for the domain $\in \mathbb{C}^n$, and range (image) $\in \mathbb{C}^m$ spaces.

Changing Bases — Rotating the Map!

Any $\vec{b} \in \mathbb{C}^m$ can be expanded in the basis of the left singular vectors of A (i.e. the columns of U), and any $\vec{x} \in \mathbb{C}^n$ in the basis of the right singular vectors of A (i.e. the columns of V)...

The coordinates for these expansions are

$$\vec{b}' = U^* \vec{b}, \quad \vec{x}' = V^* \vec{x}.$$

Now, the relation $\vec{b} = A\vec{x}$ can be written in terms of \vec{b}' and \vec{x}' :

$$\vec{b} = A\vec{x} \Leftrightarrow U^* \vec{b} = U^* A\vec{x} = U^* \underbrace{U\Sigma V^*}_A \vec{x} \Leftrightarrow \vec{b}' = \Sigma \vec{x}'$$

Singular Value vs. Eigenvalue Decomposition

1 of 2

The idea of **diagonalizing** a matrix by a change of basis is the foundation for the study of eigenvalues.

A non-defective square matrix A can be expressed as a diagonal matrix of eigenvalues Λ , if the range (image) and domain are expressed in a basis of the eigenvectors. The **eigenvalue decomposition** of $A \in \mathbb{C}^{m \times m}$ is

$$A = X\Lambda X^{-1}$$

where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_m)$, and the columns of $X \in \mathbb{C}^{m \times m}$ contain linearly independent eigenvectors of A .

We can change basis for the expression $\vec{b} = A\vec{x}$:

$$\vec{b}' = X^{-1}\vec{b}, \quad \vec{x}' = X^{-1}\vec{x}.$$

and find that

$$\vec{b}' = \Lambda\vec{x}'$$

Singular Value vs. Eigenvalue Decomposition

2 of 2

The SVD and Eigenvalue Decomposition

The SVD, $A = U\Sigma V^*$	Eigenvalue Decomp., $A = X\Lambda X^{-1}$
Properties	
Uses two different bases — the set of right and left singular vectors.	Uses one basis — the eigenvectors.
Uses orthonormal bases	Uses a basis which is generally not orthogonal.
All matrices (even rectangular ones) have a singular value decomposition.	Not all matrices (even square ones) have an eigenvalue decomposition.
(Typical) Application Relevance	
Behavior of A itself, or A^{-1} . Information in A .	Behavior of A^k , e^{tA} .

The SVD \rightsquigarrow Matrix Properties

The Rank

The SVD has many connections with other fundamental topics in linear algebra...

In the following slides, assume that $A \in \mathbb{C}^{m \times n}$, let $p = \min(m, n)$, and let $r \leq p$ denote the number of non-zero singular values of A ; finally let $\text{span}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_m)$ denote the space spanned by the vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_m$, i.e. all linear combinations of the vectors.

Theorem (Rank of a Matrix)

$$\text{rank}(A) = r.$$

Proof (Rank of a Matrix)

The rank of a diagonal matrix is the number of non-zero entries. In the decomposition $A = U\Sigma V^*$, both U and V are full rank. Therefore $\text{rank}(A) = \text{rank}(\Sigma) = r$. \square

The SVD \rightsquigarrow Matrix Properties

The Range (Image) and Null-space

Theorem (Range (Image) and Nullspace of a Matrix)

$$\begin{aligned}\text{range}(A) &= \text{span}(\vec{u}_1, \vec{u}_2, \dots, \vec{u}_r), \\ \text{null}(A) &= \text{span}(\vec{v}_{r+1}, \vec{v}_{r+2}, \dots, \vec{v}_n).\end{aligned}$$

Proof (Range (Image) and Nullspace of a Matrix)

This follows directly from the change of bases induced by $A = U\Sigma V^*$ and the fact that

$$\begin{aligned}\text{range}(\Sigma) &= \text{span}(\vec{e}_1, \vec{e}_2, \dots, \vec{e}_r) \subseteq \mathbb{C}^m, \\ \text{null}(\Sigma) &= \text{span}(\vec{e}_{r+1}, \vec{e}_{r+2}, \dots, \vec{e}_n) \subseteq \mathbb{C}^n.\end{aligned}$$

The SVD \rightsquigarrow Matrix Properties

Euclidean and Frobenius Norms

Theorem (Euclidean and Frobenius Matrix Norms)

$$\|A\|_2 = \sigma_1, \quad \text{and} \quad \|A\|_F = \sqrt{\sigma_1^2 + \sigma_2^2 + \cdots + \sigma_r^2}.$$

Proof (Euclidean and Frobenius Matrix Norms)

We already established that $\sigma_1 = \|A\|_2$ in the existence proof, since $A = U\Sigma V^*$ with unitary U and V ,

$$\|A\|_2 = \|\Sigma\|_2 = \max\{|\sigma_i|\} = \sigma_1.$$

Now, since the Frobenius norm is invariant under unitary transformations, $\|A\|_F = \|\Sigma\|_F = \sqrt{\sigma_1^2 + \sigma_2^2 + \cdots + \sigma_r^2}$.

The SVD \rightsquigarrow Matrix Properties

Singular Values / Eigenvalues

Theorem

*The non-zero singular values of A are the square roots of the non-zero eigenvalues of A^*A or AA^* (these two matrices have the same non-zero eigenvalues).*

Proof (Singular Values from AA^* or A^*A)

From

$$A^*A = (U\Sigma V^*)^*(U\Sigma V^*) = V\Sigma^*U^*U\Sigma V^* = V(\Sigma^*\Sigma)V^* = V(\Sigma^*\Sigma)V^{-1}$$

we see that A^*A and $\Sigma^*\Sigma = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_p^2)$ have the same eigenvalues, $\lambda_i = \sigma_i^2$, $i = 1, 2, \dots, p$.

If $n > p$, we have an additional $(n - p)$ zero eigenvalues.

The same argument works for AA^* (just substitute m for n)...

The SVD \rightsquigarrow Matrix Properties

Singular Values / Eigenvalues

Theorem ($\sigma_k = |\lambda_k|$ for Hermitian Matrices)

If $A = A^*$, then the singular values of A are the absolute values of the eigenvalues of A .

Note: In the language of [MATH 524] A is self-adjoint.

Proof (part 1)

The eigenvalues of a Hermitian matrix are real since if (λ, \vec{v}) is an eigenvalue-eigenvector pair ($\lambda \neq 0$), then

$$\langle \vec{v}, A\vec{v} \rangle = \vec{v}^* A\vec{v} = (A^* \vec{v})^* \vec{v} = \langle A^* \vec{v}, \vec{v} \rangle$$

$$\langle \vec{v}, A\vec{v} \rangle = \langle \vec{v}, \lambda \vec{v} \rangle = \lambda \langle \vec{v}, \vec{v} \rangle$$

$$\langle \vec{v}, A\vec{v} \rangle = \langle A^* \vec{v}, \vec{v} \rangle = \langle A\vec{v}, \vec{v} \rangle = \langle \lambda \vec{v}, \vec{v} \rangle = \lambda^* \langle \vec{v}, \vec{v} \rangle$$

Hence, $\lambda = \lambda^* \Rightarrow \lambda \in \mathbb{R}$. Further, a Hermitian matrix has a complete set of orthogonal eigenvectors. This means that we can diagonalize A

$$A = Q\Lambda Q^* = Q(|\Lambda| \text{sign}(\Lambda))Q^*$$

for some unitary matrix Q and Λ a real diagonal matrix...

The SVD \rightsquigarrow Matrix Properties

Singular Values / Eigenvalues

Proof (part 2)

Since $\text{sign}(\Lambda)Q^*$ is unitary, we have

$$A = \underbrace{Q}_U \underbrace{|\Lambda|}_\Sigma \underbrace{(\text{sign}(\Lambda)Q^*)}_{V^*}$$

a SVD of A , where $\sigma_i = |\lambda_i|$, $i = 1, 2, \dots, p$. (An appropriate ordering of the columns of U guarantees that the singular values are ordered in decreasing order.) \square

The SVD \rightsquigarrow Matrix Properties

The Determinant

Theorem

For $A \in \mathbb{C}^{m \times m}$, $|\det(A)| = \prod_{i=1}^m \sigma_i$.

Proof (Magnitude of Determinant is Product of Singular Values)

$$\begin{aligned} |\det(A)| &= |\det(U\Sigma V^*)| = |\det(U)| \cdot |\det(\Sigma)| \cdot |\det(V^*)| \\ &= 1 \cdot |\det(\Sigma)| \cdot 1 = |\det(\Sigma)| = \prod_{i=1}^m \sigma_i \end{aligned}$$

where we have used the fact that $\det(AB) = \det(A)\det(B)$ and that the magnitude of the determinant of a unitary matrix is one.

The SVD \rightsquigarrow Matrix Properties

Low-Rank Approximations, 1 of 5

This discussion is a significant part of WHY this course exists!

Given the SVD of A , $A = U\Sigma V^*$, we can represent A as a sum of r rank-one matrices

$$A = \sum_{k=1}^r \sigma_k \vec{u}_k \vec{v}_k^*$$

This is certainly not the only way to write A as a sum of rank-one matrices: it could be written as a sum of its m rows, n columns, or even its mn entries...

The decomposition above has the special property that if we truncate the sum at $\nu < r$, then that partial sum captures as much "energy" of A as possible for a rank- ν sub-matrix of A .

We formalize this in a theorem...

Theorem (Optimal Low-Rank Approximation)

For any ν with $0 \leq \nu < r$, define

$$A_\nu = \sum_{k=1}^{\nu} \sigma_k \vec{u}_k \vec{v}_k^*$$

if $\nu = p = \min(m, n)$, define $\sigma_{\nu+1} = 0$. Then

$$\|A - A_\nu\|_2 = \inf_{\substack{B \in \mathbb{C}^{m \times n} \\ \text{rank}(B) \leq \nu}} \|A - B\|_2 = \sigma_{\nu+1}$$

The SVD \rightsquigarrow Matrix Properties

Low-Rank Approximations, 2.5 of 5

Low Rank Approximations in DS/Machine Learning/Generative AI

Low-Rank Adaptation (LoRA) is a family of methods for fine-tuning large-scale AI/Machine Learning models in an efficient manner.

"Base-Models" (e.g. LLMs like ChatGPT; or image-generative models like the Stable Diffusion SD1.5 or SDXL models) are trained on extremely large data sets — this training uses significant resources, *i.e.* they are "expensive."

Very Simplified: fine-tuning is "retraining" (parts of) the model using a smaller specific data set; e.g. published peer-reviewed mathematics research papers, or images created in a particular "style."

The Model parameters use usually collected in a large matrix $A \in \mathbb{R}^{M \times N}$; and the fine-tuning computes "a few" — collected in much smaller matrices $B \in \mathbb{R}^{M \times p}$, and $C \in \mathbb{R}^{p \times N}$, so that the effective fine-tuned model can be represented as

$$A + BC$$

M and N are usually "quite large" ($> 1,000$), and p "small" (< 10).

The SVD \rightsquigarrow Matrix Properties

Low-Rank Approximations, 3 of 5

Proof (Optimal Low-Rank Approximation)

Suppose that there is some B with $\text{rank}(B) \leq \nu$ such that $\|A - B\|_2 < \|A - A_\nu\|_2 = \sigma_{\nu+1}$.

Then there is an $(n - \nu)$ -dimensional subspace $\text{null}(B) = \mathbb{W} \subseteq \mathbb{C}^n$ such that $\vec{w} \in \mathbb{W} \Rightarrow B\vec{w} = 0$. Thus $\forall \vec{w} \in \mathbb{W}$:

$$\|A\vec{w}\|_2 = \|(A - B)\vec{w}\|_2 \leq \|A - B\|_2 \|\vec{w}\|_2 < \sigma_{\nu+1} \|\vec{w}\|_2.$$

Now, \mathbb{W} is an $(n - \nu)$ -dimensional subspace where $\|A\vec{w}\|_2 < \sigma_{\nu+1} \|\vec{w}\|_2$. But there is a $(\nu + 1)$ -dimensional subspace where $\|A\vec{w}\|_2 \geq \sigma_{\nu+1} \|\vec{w}\|_2$ — $\mathbb{V} = \text{span}(u_1, \dots, u_{\nu+1})$ the space spanned by the first $(\nu + 1)$ right singular vectors of A .

Since the sum of the dimensions of the two subspaces $(\nu + 1) + (n - \nu) = (n + 1)$ exceeds n , there must be a non-zero vector lying in both. This is a contradiction.

The SVD \rightsquigarrow Matrix Properties

Low-Rank Approximations, 4 of 5

The preceding theorem has a nice geometrical interpretation.

Ponder the issue of finding the best approximation of an n -dimensional hyper-ellipsoid.

- \Rightarrow The best approximation by a 2-dimensional ellipse must be the ellipse spanned by the largest and second largest axis.
- \Rightarrow We get the best 3-dimensional approximation by adding the span of the 3rd largest axis, etc...

This is useful in many applications, *e.g.* signal compression (images, audio, etc.), analysis of large data sets, etc.

The SVD \rightsquigarrow Matrix Properties

Low-Rank Approximations, 5 of 5

We state the following theorem, and leave the proof as an "exercise."

Theorem

For the matrix A_ν as defined in the previous theorem

$$\|A - A_\nu\|_F = \inf_{\substack{B \in \mathbb{C}^{m \times n} \\ \text{rank}(B) \leq \nu}} \|A - B\|_F = \sqrt{\sigma_{\nu+1}^2 + \sigma_{\nu+2}^2 + \cdots + \sigma_r^2}$$

We will get back to **how to compute** the SVD later. For now, we note that it is a powerful tool which can be used to

- find the numerical rank of a matrix;
- find the orthonormal basis for the range (image) and null-space;
- computing $\|A\|_2$;
- computing low-rank approximations.

The SVD shows up in least squares fitting, regularization, intersection of subspaces (video games?), and many, many other problems.