

# Numerical Matrix Analysis

## Notes #25 Arnoldi Iteration

Peter Blomgren  
([blomgren@sdsu.edu](mailto:blomgren@sdsu.edu))

Department of Mathematics and Statistics  
Dynamical Systems Group  
Computational Sciences Research Center  
San Diego State University  
San Diego, CA 92182-7720

<http://terminus.sdsu.edu/>

Spring 2024  
(Revised: April 24, 2024)



## Outline

- 1 **Arnoldi Iteration**
  - Introduction
  - Mechanics
  - Interpretations
  
- 2 **Arnoldi Iteration  $\rightsquigarrow$  Eigenvalues**
  - Computing Eigenvalues by Arnoldi Iteration
  - The Arnoldi/Lanczos Approximation Problem
  - Convergence

## Introduction, Definitions

The Arnoldi Iteration generates a **sequence of matrices** from which the eigenvalues  $\lambda$  of  $A\vec{x} = \lambda\vec{x}$  for  $A \neq A^*$  can easily be computed.

It is a Gram-Schmidt-style orthogonalization iteration for transforming the matrix into Hessenberg form.

	$A \rightsquigarrow QR$	$A \rightsquigarrow QHQ^*$
Orthogonal "Structuring" 'Structured' Orthogonalization	Householder Gram-Schmidt	Householder Arnoldi

## Definition (Krylov Sequences and Subspaces)

Given a vector  $\vec{x}$  and a matrix  $A$ , the associated **Krylov vector sequence** is  $\{\vec{x}, A\vec{x}, \dots, A^{k-1}\vec{x}, \dots\}$ , and the corresponding **Krylov subspaces**  $K(A, \vec{x}; k) = \text{span}(\vec{x}, A\vec{x}, \dots, A^{k-1}\vec{x})$ .

## Householder vs. Gram-Schmidt/Arnoldi

In the context of the QR factorization we have the Householder reflections which “triangularize”  $A$  by a sequence of orthogonal operations; and the Gram-Schmidt process which Orthogonalize  $A$  by a sequence of triangular operations.

Row-  
Ops.

The Householder reflection strategy performs better (in terms of  $Q$ -orthonormality) in finite precision.

Col-  
Ops.

The Gram-Schmidt process has the **advantage that it can be stopped part-way**; yielding an incomplete QR-factorization of the first  $n$  columns of  $A$ .

We have viewed the process of computing the Hessenberg (upper triangular + 1 sub-diagonal) form  $A = QHQ^*$  as a two-sided application of Householder reflections. The Arnoldi iteration is the “interruptible” Gram-Schmidt “equivalent.”

Hessenberg Reduction, at step  $n$ 

1 of 3

## Problem Structure and Notation

We consider the problem of computing  $A \rightsquigarrow QHQ^*$  for  $A \in \mathbb{R}^{m \times m}$ . All norms  $\|\cdot\|$  are 2-norms  $\|\cdot\|_2$ .  $n$  is the iteration number, and we operate in the regime where  $n \ll m$ .

The complete Hessenberg reduction can be written in the form  $A = QHQ^*$ , or  $AQ = QH$ , where  $A, Q, H \in \mathbb{R}^{m \times m}$ . However, seen at the  $n^{\text{th}}$  iterative step, we consider the partially computed matrix  $Q_n \in \mathbb{R}^{m \times n}$ , *i.e.* the first  $n$  columns of  $Q$ :

$$Q_n = \begin{bmatrix} | & | & & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{bmatrix}.$$

Hessenberg Reduction, at step  $n$ 

2 of 3

We also consider the upper  $((n+1) \times n)$  block of  $H$ , also a Hessenberg matrix

$$\tilde{H}_n = \begin{bmatrix} h_{1,1} & & \cdots & & h_{1,n} \\ h_{2,1} & h_{2,2} & & & \vdots \\ & \ddots & \ddots & & \\ & & & h_{n,n-1} & h_{n,n} \\ & & & & h_{n+1,n} \end{bmatrix},$$

so that

$$AQ_n = Q_{n+1}\tilde{H}_n$$

is well defined, since

$$[m \times n] = [m \times m] \otimes [m \times n] \implies [m \times (n+1)] \otimes [(n+1) \times n] = [m \times n].$$



Hessenberg Reduction, at step  $n$ 

3 of 3

The  $n^{\text{th}}$  column of the relation  $AQ_n = Q_{n+1}\tilde{H}_n$  is

$$A\vec{q}_n = \sum_{k=1}^{n+1} h_{k,n}\vec{q}_k,$$

which means that  $\vec{q}_{n+1}$  satisfies an  $(n+1)$ -term recurrence relation involving itself, and previous Krylov basis vectors,  $\vec{q}_1, \dots, \vec{q}_n$ :

$$\vec{q}_{n+1} = \frac{1}{h_{n+1,n}} \left[ A\vec{q}_n - \sum_{k=1}^n h_{k,n}\vec{q}_k \right].$$

The Arnoldi iteration implements this recurrence relation.



## The Arnoldi Algorithm

## Algorithm (Arnoldi Iteration)

```
1:  $\vec{b} \leftarrow \text{random}(\mathbb{R}^{m \times 1})$ ,  
2:  $\vec{q}_1 \leftarrow \vec{b} / \|\vec{b}\|$   
3: for  $n \in \{1, 2, \dots\}$  do  
4:    $\vec{v} \leftarrow A\vec{q}_n$   
5:   for  $j \in \{1, \dots, n\}$  do  
6:      $h_{j,n} \leftarrow \vec{q}_j^* \vec{v}$   
7:      $\vec{v} \leftarrow \vec{v} - h_{j,n}\vec{q}_j$   
8:   end for  
9:    $h_{n+1,n} \leftarrow \|\vec{v}\|$   
10:   $\vec{q}_{n+1} \leftarrow \vec{v} / h_{n+1,n}$   
11: end for
```

TB-33.2:  $h_{n+1,n} = 0$  (Breakdown due to Convergence)



The power of the Arnoldi process is the various interpretations that can be made of it (and its by-products), and the algorithms these suggest.

### Arnoldi: Generated Krylov Subspaces

From construction, the generated vectors  $\{\vec{q}_k\}$  form orthonormal bases for the Krylov subspaces generated by  $A$  and  $\vec{b}$ :

$$K(A, \vec{b}; k) = \text{span}(\vec{b}, A\vec{b}, \dots, A^{k-1}\vec{b}) = \text{span}(\vec{q}_1, \vec{q}_2, \dots, \vec{q}_k) \subseteq \mathbb{C}^m.$$

The  $(m \times n)$  Krylov matrix  $\mathcal{K}_n$  must have a reduced  $QR$ -factorization:

$$\mathcal{K}_n = \left[ \begin{array}{c|c|c|c} \vec{b} & A\vec{b} & \dots & A^{n-1}\vec{b} \end{array} \right] = Q_n R_n,$$

where  $Q_n$  is the matrix previously defined ( $\vec{q}_k$  are its columns).

Neither  $\mathcal{K}_n$ , nor  $R_n$  can be stably formed during the Arnoldi process, but they hint at why the Arnoldi process leads to effective methods for determining certain (the dominant) eigenvalues of  $A$ .

The relationship between the Arnoldi iteration, and direct QR factorization of  $\mathcal{K}_n$  is reminiscent of the Simultaneous-(Power)-Iteration vs. QR-algorithm approaches:

*Straight-forward, unstable*

Simultaneous Iteration  
 $\mathcal{K}_n = Q_n R_n$

*Subtle, stable*

QR-algorithm  
Arnoldi

We can also view the Arnoldi process as a computation of **projections onto successive Krylov subspaces**.

Note:  $(n \times (n + 1))$  Identity

The product

$$Q_n^* Q_{n+1} = I = \delta_{k,\ell}, \quad k = 1, \dots, n, \quad \ell = 1, \dots, (n + 1).$$

Therefore  $Q_n^* Q_{n+1} \tilde{H}_n$  is the  $(n \times n)$  Hessenberg matrix obtained by removing the last row of  $\tilde{H}_n$ :

$$H_n = \begin{bmatrix} h_{1,1} & & \cdots & & h_{1,n} \\ h_{2,1} & h_{2,2} & & & \vdots \\ & \ddots & \ddots & & \\ & & & h_{n,n-1} & h_{n,n} \end{bmatrix}, \quad \text{and } H_n = Q_n^* A Q_n.$$

This matrix can be interpreted as the orthogonal projection ( $P : \mathbb{R}^{m \times m} \mapsto \mathbb{R}^{m \times n}$ ) of  $A$  onto  $\mathcal{K}_n$ , represented by the basis  $\{\vec{q}_1, \dots, \vec{q}_n\}$ . (Exact statement in TB pp.254, it makes sense if you have taken [MATH 524]).

This type of projection shows up in a variety of contexts, and is sometimes referred to as the *Rayleigh-Ritz* procedure. It turns out that the diagonal elements  $H_n$  are the Rayleigh coefficients of  $A$  with respect to the vectors  $\vec{q}_j$ .

This projection process is one of the ideas underlying the *Finite Element Method* for solution of PDEs, as well as *spectral methods*.

Since  $H_n$  is a projection of  $A$ , its eigenvalues  $\lambda(H_n)$  are related to the eigenvalues of  $A$ ; they are referred to as the *Arnoldi eigenvalue estimates*, or *Ritz values* (with respect to  $\mathcal{K}_n$ ) of  $A$ .



## Arnoldi: Summary (so far)

## Theorem

The matrices  $Q_n$  generated by the Arnoldi iteration are reduced QR-factors of the Krylov matrix:

$$\mathcal{K}_n = Q_n R_n.$$

The Hessenberg matrices  $H_n$  are the corresponding projections

$$H_n = Q_n^* A Q_n,$$

and the successive iterates are related by

$$A Q_n = Q_{n+1} \tilde{H}_n.$$

## Arnoldi Eigenvalue Computation Strategy

## Strategy

- 1 Perform Arnoldi Iteration
- 2 Compute the Eigenvalues of  $H_n$  using e.g. the QR-algorithm at regular intervals.
  - $\lambda(H_n)$  are the *Arnoldi eigenvalue estimates*, or *Ritz values*.
  - a (growing) subset of  $\lambda(H_n)$  typically converge quickly, and are eigenvalues of  $A$ .
  - [∃ Movie]
  - Typically, the largest-modulus eigenvalues are located first; and usually (but not always) those are the ones we are interested in.

Some aspects of the convergence can be quantified...

## Minimization Problem Over the Monic Polynomials

Let  $\vec{x} \in K(A, \vec{b}; n)$ , then

$$\vec{x} = \sum_{k=0}^{n-1} c_k A^k \vec{b} = q(A) \vec{b},$$

where  $q(A)$  is a (matrix) polynomial in  $A$ .

With

$$\mathcal{P}^n = \{ \text{ALL MONIC POLYNOMIALS OF DEGREE } n \},$$

where *monic* means  $c_n = 1$ , we can state the

## Arnoldi (and Lanczos) Approximation Problem (AAP)

Find  $p_n \in \mathcal{P}^n$  such that

$$\|p_n(A)\vec{b}\| \text{ is minimized.}$$



## Solution to the Minimization Problem

The Arnoldi iteration solves the stated minimization problem:

## Theorem

As long as the Arnoldi iteration does not break down (i.e.  $K_n$  is of full rank  $n$ ), (AAP) has a unique solution  $p_n$ ; ~~the characteristic~~ *minimal polynomial* of  $H_n$ . [Proof in Trefethen & Bau, pp. 259–260.]

Hence, the Ritz values generated by the Arnoldi iteration are the roots of the optimal polynomial.

Since the class of monic polynomials  $\mathcal{P}^n$  is invariant with respect to translations  $[z \mapsto (z + \alpha)]$ , so is the Arnoldi iteration:

**Ref:** See [MATH 524 (NOTES#8)] for discussions on Cayley–Hamilton Theorem; the Minimal, and Characteristic Polynomials.



## Invariance Theorem

## Theorem

Let the Arnoldi iteration be applied to a matrix  $A \in \mathbb{C}^{m \times m}$ ; each of the following invariance properties hold:

- **[TRANSLATION-INVARIANCE]** If  $A$  is changed to  $(A + \sigma I)$  for some  $\sigma \in \mathbb{C}$ , and  $\vec{b}$  is left unchanged, the Ritz values  $\{\theta_j\}$  change to  $\{\theta_j + \sigma\}$ .
- **[SCALE-INVARIANCE]** If  $A$  is changed to  $\sigma A$  for some  $\sigma \in \mathbb{C}$ , and  $\vec{b}$  is left unchanged, the Ritz values  $\{\theta_j\}$  change to  $\{\sigma\theta_j\}$ .
- **[INVARIANCE UNDER UNITARY SIMILARITY TRANSFORMS]** If  $A$  is changed to  $UAU^*$  for some unitary matrix  $U$ , and  $\vec{b}$  is changed to  $U\vec{b}$ , the Ritz values do not change.

In all three cases the Ritz vectors, the vectors  $Q_n \vec{y}_j$  corresponding to the eigenvectors  $\vec{y}_j$  of  $H_n$ , do not change.



## Invariance Theorem: Discussion

We know (Schur Factorization) that every matrix is similar to an upper triangular matrix.

Therefore, the invariance property shows that we can understand the properties of the Arnoldi iteration by considering upper triangular matrices.

Note: NON-HERMITIAN MATRICES  $\leftrightarrow$  TRIANGULAR MATRICES, and HERMITIAN MATRICES  $\leftrightarrow$  DIAGONAL MATRICES. Hence in the non-Hermitian case off-diagonal elements complete the description of the matrix action beyond “just” the eigenvalues.

# Convergence

**Easy Case:**  $A$  is diagonalizable, and has  $n \ll m$  distinct eigenvalues. In this case the Arnoldi iteration finds the  $n$  eigenvalues exactly, as long as the “seed vector”  $\vec{b}$  has non-zero components in each eigenspace  $E_{\lambda_k}$ ; and  $\|p_n(A)\vec{b}\| = 0$ . [In infinite precision]

**General Case:** Here, the Ritz values are approximations to eigenvalues, and  $p_n$  is a *pseudo-minimal* polynomial; i.e.  $\|p_n(A)\vec{b}\|$  is small.

The convergence process can be illustrated by a particular level-curve (*lemniscate*) of the polynomial.

## Arnoldi Lemniscates

## Definition (Lemniscate, Level Set)

A lemniscate is a curve, or collection of curves

$$\{z \in \mathbb{C} : |p(z)| = \mathcal{C}\},$$

where  $p(z)$  is a polynomial and  $\mathcal{C}$  is a real constant.

If  $p(z)$  is the Arnoldi polynomial  $p_n(\cdot)$  for an Arnoldi iteration, and

$$\mathcal{C}_n = \frac{\|p_n(A)\vec{b}\|}{\|\vec{b}\|},$$


then the curves are called *Arnoldi Lemniscates*.

The components of the *Arnoldi Lemniscates* tend to surround the extreme eigenvalues of  $A$  and the shrink rapidly to a point (the eigenvalue).


# Arnoldi Lemniscates

1 & 2 of 16

Arnoldi Eigenvalue Estimates 1 / 128




Arnoldi Eigenvalue Estimates 2 / 128




# Arnoldi Lemniscates

3 & 4 of 16

Arnoldi Eigenvalue Estimates 3 / 128




Arnoldi Eigenvalue Estimates 4 / 128




# Arnoldi Lemniscates

5 & 6 of 16

Arnoldi Eigenvalue Estimates 5 / 128



Arnoldi Eigenvalue Estimates 6 / 128







# Arnoldi Lemniscates

7 & 8 of 16

Arnoldi Eigenvalue Estimates 7 / 128




Arnoldi Eigenvalue Estimates 8 / 128




# Arnoldi Lemniscates

9 & 10 of 16

Arnoldi Eigenvalue Estimates 9 / 128




Arnoldi Eigenvalue Estimates 10 / 128




# Arnoldi Lemniscates

11 & 12 of 16

Arnoldi Eigenvalue Estimates 11 / 128




Arnoldi Eigenvalue Estimates 12 / 128




# Arnoldi Lemniscates

13 & 14 of 16

Arnoldi Eigenvalue Estimates 13 / 128




Arnoldi Eigenvalue Estimates 14 / 128




# Arnoldi Lemniscates

15 & 16 of 16

Arnoldi Eigenvalue Estimates 15 / 128



Arnoldi Eigenvalue Estimates 16 / 128



## Rate of Convergence

The rate of convergence is not well understood in general.

For a single outlying eigenvalue, the convergence rate is typically geometric (in the first few iterations, then faster); e.g.

Trefethen & Bau show an example where

$$|\lambda^{(n)} - \lambda| \sim \left(\frac{2}{3}\right)^n.$$

Convergence is highly dependent on the shape of the eigenvalue spectrum  $\lambda(A)$ . — In the next lecture we will see some examples of this in the context of the related GMRES algorithm.