

Numerical Matrix Analysis

Notes #4

Matrix Norms, the Singular Value Decomposition

Peter Blomgren

`<blomgren@sdsu.edu>`

Department of Mathematics and Statistics

Dynamical Systems Group

Computational Sciences Research Center

San Diego State University

San Diego, CA 92182-7720

<http://terminus.sdsu.edu/>

Spring 2024

(Revised: February 20, 2024)



Outline

- 1 Student Learning Targets, and Objectives
 - SLOs: Matrix Norms, and the Singular Value Decomposition
- 2 Introduction: Matrix Norms
 - Recap
 - Inequalities
 - General Matrix Norms
- 3 The Singular Value Decomposition
 - Many Names... One Powerful Tool!
 - Examples for 2×2 Matrices
 - More Details, and Examples Revisited
- 4 The SVD of a Matrix: Formal Definition
 - Spheres and Hyper-ellipses
 - Homework

Student Learning Targets, and Objectives

Target Matrix Norms

Objective Special Cases: 1- and ∞ -norms

Objective Hölder, and Cauchy-Bunyakovsky-Schwarz inequality

Objective Special Case: 2-norm of rank-1 matrices

Objective Frobenius (Hilber-Schmidt) norm

Target The Singular Value Decomposition (the SVD)

Objective In the first pass: the SVD as a concept, and geometrical interpretation

Objective Fundamental language and concepts:

- principal semi-axes
- Singular values (σ_k , Σ)
- Left singular vectors (\vec{u}_k , U)
- Right singular vectors (\vec{v}_k , V)

Last Time

Orthogonal Vectors, Matrices and Norms:

- The Adjoint / Hermitian Conjugate of a Matrix, A^*
- The Inner Product of Two Vectors, $\langle \vec{x}, \vec{y} \rangle = \vec{x}^* \vec{y}$
- Orthogonal, $\langle \vec{x}, \vec{y} \rangle = 0$, and Orthonormal, $\|\vec{x}\| = 1$, Vectors
- Orthogonal and Orthonormal Sets — Linear Independence; Basis for \mathbb{C}^m
- Unitary Matrices $Q^* Q = I$
- Vector Norms, $\|\cdot\|_p$ (p -norms), weighted p -norms
- Induced Matrix Norms

Inequalities: Hölder and Cauchy-Bunyakovsky-Schwarz

1 of 2

Last time we noted that the 1-norm and ∞ -norm of a matrix simplify to the maximal column- and row-sum, respectively, *i.e.* for $A \in \mathbb{C}^{m \times n}$

$$\|A\|_1 = \max_{1 \leq j \leq n} \|\vec{a}_j\|_1$$

$$\|A\|_\infty = \max_{1 \leq i \leq m} \|\vec{a}_i^*\|_1$$

For other p -norms, $1 \leq p \leq \infty$, the matrix-norms do not reduce to simple direct-computable expressions like the ones above.

Inequalities: Hölder and Cauchy-Bunyakovsky-Schwarz

2 of 2

However, we can usually find useful bounds on vector- and matrix-norms, using the **Hölder inequality**:

$$|\vec{x}^* \vec{y}| \leq \|\vec{x}\|_p \|\vec{y}\|_q, \quad \frac{1}{p} + \frac{1}{q} = 1.$$

In the special case $p = q = 2$, the inequality is known as the **Cauchy-Schwarz**, or **Cauchy-Bunyakovsky-Schwarz** inequality

$$|\vec{x}^* \vec{y}| \leq \|\vec{x}\|_2 \|\vec{y}\|_2.$$

Augustin-Louis Cauchy, 21 August 1789 – 23 May 1857. (French)

⇒ proof for sums (1821).

Viktor Yakovlevich Bunyakovsky, 16 December 1804 – 12 December 1889. (Russian, Cauchy's graduate student)

⇒ proof for integrals (1859).

Karl Hermann Amandus Schwarz, 25 January 1843 – 30 November 1921. (German)

⇒ Modern proof (1888).

Otto Ludwig Hölder, 22 December 1859 – 29 August 1937

Example: 2-norm of a Rank-1 Matrix $A = \vec{u}\vec{v}^*$

1 of 2

Rank-1 matrices formed by an outer product $\vec{u}\vec{v}^*$ show up in many numerical schemes:

$$\vec{u}\vec{v}^* = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_m \end{bmatrix} \begin{bmatrix} v_1^* & v_2^* & \cdots & v_n^* \end{bmatrix} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_m \end{bmatrix} \vec{v}^* = \begin{bmatrix} u_1 \vec{v}^* \\ u_2 \vec{v}^* \\ \vdots \\ u_m \vec{v}^* \end{bmatrix}$$

Now, for any $\vec{x} \in \mathbb{C}^n$, we get

$$\|A\vec{x}\|_2 = \|\vec{u}\vec{v}^*\vec{x}\|_2 = \|\vec{u}\|_2 |\vec{v}^*\vec{x}| \leq \|\vec{u}\|_2 \|\vec{v}\|_2 \|\vec{x}\|_2$$

Hence,

$$\|A\|_2 = \sup_{\vec{x} \in \mathbb{C}^n - \{\vec{0}\}} \frac{\|A\vec{x}\|_2}{\|\vec{x}\|_2} \leq \|\vec{u}\|_2 \|\vec{v}\|_2$$

Example: 2-norm of a Rank-1 Matrix $A = \vec{u}\vec{v}^*$

2 of 2

Since $\vec{v} \in \mathbb{C}^n$, the inequality

$$\|A\|_2 = \sup_{\vec{x} \in \mathbb{C}^n - \{\vec{0}\}} \frac{\|A\vec{x}\|_2}{\|\vec{x}\|_2} \leq \|\vec{u}\|_2 \|\vec{v}\|_2$$

is actually an equality. Let $\vec{x} = \vec{v}$:

$$\|A\vec{v}\|_2 = \|\vec{u}\vec{v}^*\vec{v}\|_2 = \|\vec{u}\|_2 |\vec{v}^*\vec{v}| = \|\vec{u}\|_2 \|\vec{v}\|_2^2$$

Bounds on the Norms of Matrix Products, $\|AB\|$

Let $A \in \mathbb{C}^{\ell \times m}$, $B \in \mathbb{C}^{m \times n}$, and $\vec{x} \in \mathbb{C}^n$: and let $\|\cdot\|$ denote compatible p -norms, then

$$\|AB\vec{x}\| \leq \|A\| \|B\vec{x}\| \leq \|A\| \|B\| \|\vec{x}\|.$$

Therefore, we have

$$\|AB\| \leq \|A\| \|B\|,$$

where, in general $\|AB\| \neq \|A\| \|B\|$.

General (Non-Induced) Matrix Norms

Matrix norms induced by vector norms are quite common, but as long as the following norm-conditions are satisfied:

- (1) $\|A\| \geq 0$, and $\|A\| = 0$ only if $A = 0$
- (2) $\|A + B\| \leq \|A\| + \|B\|$
- (3) $\|\alpha A\| = |\alpha| \|A\|$

for $A \in \mathbb{C}^{m \times n}$, then $\|\cdot\|$ is a valid matrix-norm.

The most commonly used non-induced matrix norm is the **Frobenius norm** (sometimes referred to as the **Hilbert-Schmidt norm**):

$$\|A\|_F = \left[\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2 \right]^{1/2}.$$

The Frobenius Norm

We can view the Frobenius Norm in terms of column- or row-sums:

$$\|A\|_F = \left[\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2 \right]^{1/2} = \left[\sum_{j=1}^n \|\vec{a}_j\|_2^2 \right]^{1/2} = \left[\sum_{i=1}^m \|\vec{a}_i^*\|_2^2 \right]^{1/2}$$

...or in terms of the trace (sum of diagonal entries)

$$\|A\|_F = \sqrt{\text{trace}(A^*A)} = \sqrt{\text{trace}(AA^*)}$$

Invariance under Unitary Multiplication

Both the 2-norm and the Frobenius norm are invariant under multiplication by unitary matrices, *i.e.*

Theorem

For any $A \in \mathbb{C}^{m \times n}$ and unitary $Q \in \mathbb{C}^{m \times m}$, we have

$$\|QA\|_2 = \|A\|_2, \quad \|QA\|_F = \|A\|_F$$

... an indication of the importance (and usefulness) of unitary matrices!

This ends our quick introduction to basic linear algebra concepts

Next: A first look at the Singular Value Decomposition (SVD)

Introduction to Linear Algebra — “Optional” for Math 254

Gilbert Strang, Wellesley-Cambridge Press, 5th edition (2016), ISBN-0980232775

Linear Algebra Done Right — “Required” for Math 524

Sheldon Axler, Springer-Verlag, 4th edition (2024), ISBN-978-3-031-41026-0, OPEN ACCESS, PDF Available:

<https://doi.org/10.1007/978-3-031-41026-0>, OR

<https://link.springer.com/content/pdf/10.1007/978-3-031-41026-0.pdf>

Linear Algebra and Learning from Data

Gilbert Strang, Wellesley-Cambridge Press, 1st edition (January 2, 2019), ISBN-0692196382

The Singular Value Decomposition

The SVD [*mathematics*] is known by many names:

- Proper Orthogonal Decomposition (POD)
- Karhunen-Loève (KL-) Decomposition [*signal analysis*]
- Principal Component Analysis (PCA) [*statistics*]
- Empirical Orthogonal Functions, etc...

“[The SVD is] absolutely a high point of linear algebra.”
Prof. Gilbert Strang, MIT

Hits on scholar.google.com




Figure: The many names, faces, and close relatives of the Singular Value Decomposition... Number of hits for "Proper.Orthogonal.Decomposition", "Empirical.Orthogonal.(Function|Functions)", "Karhunen.Loeve", "Canonical.Correlation.Analysis"

Hits on scholar.google.com




Figure: The many names, faces, and close relatives of the Singular Value Decomposition... Number of hits for "Proper.Orthogonal.Decomposition", "Empirical.Orthogonal.(Function|Functions)", "Karhunen.Loeve", "Canonical.Correlation.Analysis", "Singular.Value.Decomposition", "Principal.Component.Analysis"

Hits on scholar.google.com




Figure: The many names, faces, and close relatives of the Singular Value Decomposition... Number of hits for "Proper.Orthogonal.Decomposition", "Empirical.Orthogonal.(Function|Functions)", "Karhunen.Loeve", "Canonical.Correlation.Analysis", "Singular.Value.Decomposition", "Principal.Component.Analysis"

The Singular Value Decomposition

In our first look at the SVD, we will **not** consider **how to** compute the SVD, but will focus on the meaning of the SVD; — especially its geometric interpretation.

The motivating geometric fact:

The image of the unit sphere under any $(m \times n)$ matrix, A , is a hyper-ellipse.

The hyper-ellipse in \mathbb{R}^m is the surface we get when stretching the unit sphere by some factors $\sigma_1, \sigma_2, \dots, \sigma_m$ in some orthogonal directions $\vec{u}_1, \vec{u}_2, \dots, \vec{u}_m$.

We take \vec{u}_i to be unit vectors, *i.e.* $\|\vec{u}_i\|_2 = 1$, thus the vectors $\{\sigma_i \vec{u}_i\}$ are the **principal semi-axes** of the hyper-ellipse.

The Singular Value Decomposition

For $A \in \mathbb{R}^{m \times n}$, if $\text{rank}(A) = r$, then exactly r of the lengths σ_i will be non-zero. In particular, if $m \geq n$, at most n of them will be non-zero.


Before we take this discussion further, let's look at some examples of the SVD of some (2×2) matrices.

Keep in mind that computing the SVD of a matrix A answers the question:

“What are the principal semi-axes of the hyper-ellipse generated when A operates on the unit sphere?”

In some sense, this constitutes to most complete information you can extract from a matrix.


Example#1: SVD of a 2×2 Matrix



$$\text{SVD} \left(\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \right) = \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}}_{[\vec{u}_1 \quad \vec{u}_2]} \underbrace{\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}}_{\text{diag}(\sigma_1, \sigma_2)} \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^*}_{[\vec{v}_1 \quad \vec{v}_2]^*}$$

For now, let's sweep the matrix V^* under the carpet, and note that the SVD has identified the directions of stretching (\vec{u}_1, \vec{u}_2) and the amount of stretching $(\sigma_1, \sigma_2) = (2, 1)$.

Example#2: SVD of a 2×2 Matrix




$$\text{SVD} \left(\begin{bmatrix} 1.6 & -0.65 \\ -0.65 & -0.3 \end{bmatrix} \right) = \begin{bmatrix} -0.9553 & 0.2955 \\ 0.2955 & 0.9553 \end{bmatrix} \begin{bmatrix} 1.8011 & 0 \\ 0 & 0.5011 \end{bmatrix} \begin{bmatrix} -0.9553 & -0.2955 \\ 0.2955 & -0.9553 \end{bmatrix}^*$$

Here, the principal semi-axes of the ellipse are

$$\sigma_1 \vec{u}_1 = 1.8011 \begin{bmatrix} -0.9553 \\ 0.2955 \end{bmatrix}, \quad \sigma_2 \vec{u}_2 = 0.5011 \begin{bmatrix} 0.2955 \\ 0.9553 \end{bmatrix}$$

Example#3: SVD of a 2×2 Matrix



$$\text{SVD} \left(\begin{bmatrix} 1.4 & -0.62 \\ -1.1 & -1.7 \end{bmatrix} \right) = \begin{bmatrix} -0.2501 & 0.9682 \\ 0.9682 & 0.2501 \end{bmatrix} \begin{bmatrix} 2.0556 & 0 \\ 0 & 1.4896 \end{bmatrix} \begin{bmatrix} -0.6885 & 0.7253 \\ -0.7253 & -0.6885 \end{bmatrix}^*$$

Here, the principal semi-axes of the ellipse are

$$\sigma_1 \vec{u}_1 = 2.0556 \begin{bmatrix} -0.2501 \\ 0.9682 \end{bmatrix}, \quad \sigma_2 \vec{u}_2 = 1.4896 \begin{bmatrix} 0.9682 \\ 0.2501 \end{bmatrix}$$

The Singular Value Decomposition

More Details, 1 of 2

Let \mathbb{S}^{n-1} be the unit sphere in \mathbb{R}^n , *i.e.*

$$\mathbb{S}^{n-1} = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\|_2 = 1\}$$

Let $A \in \mathbb{R}^{m \times n}$ ($m \geq n$) be of full rank, *i.e.* $\text{rank}(A) = n$, and let $A\mathbb{S}^{n-1}$ denote the image of the unit sphere (our hyper-ellipse).

The n **singular values of A** are the lengths of the n principal semi-axes of $A\mathbb{S}^{n-1}$ (some lengths may be zero), written as $\sigma_1, \sigma_2, \dots, \sigma_n$. By convention, they are ordered in descending order, so that

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n > 0$$

The n **left singular vectors of A** are the unit vectors $\{\vec{u}_1, \vec{u}_2, \dots, \vec{u}_n\}$ oriented in the directions of the principal semi-axes of $A\mathbb{S}^{n-1}$.

The Singular Value Decomposition

More Details, 2 of 2

Note that the vector $\sigma_k \vec{u}_k$ is the k th largest principal semi-axis of AS^{n-1} .

The n **right singular vectors of A** are the unit vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\} \in \mathbb{S}^{n-1}$ that are **pre-images** of the principal semi-axes of AS^{n-1} , *i.e.*

$$A\vec{v}_k = \sigma_k \vec{u}_k.$$


Note how this is similar to *and* different from an eigen-vector – eigen-value pair:

$$A\vec{\xi}_k = \lambda_k \vec{\xi}_k.$$


With that knowledge we can re-visit the three examples...

Revisited: Our 3 Examples


3D Movies \exists






$$A = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$



$$A = \begin{bmatrix} 1.6 & -0.65 \\ -0.65 & -0.3 \end{bmatrix}$$



$$A = \begin{bmatrix} 1.4 & -0.62 \\ -1.1 & -1.7 \end{bmatrix}$$



The Reduced SVD

1 of 2

What we have described so far is known as the reduced (or thin) SVD, if we collect the relations between the right and left singular vectors,

$$A\vec{v}_k = \sigma_k \vec{u}_k, \quad k = 1, \dots, n$$

in full-blown matrix notation we get

$$\begin{bmatrix} A \end{bmatrix} \begin{bmatrix} \left| \vec{v}_1 \right| & \left| \vec{v}_2 \right| & \cdots & \left| \vec{v}_n \right| \end{bmatrix} = \begin{bmatrix} \left| \vec{u}_1 \right| & \left| \vec{u}_2 \right| & \cdots & \left| \vec{u}_n \right| \end{bmatrix} \begin{bmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \end{bmatrix}$$

Usually written in the compact form

$$AV = \hat{U}\hat{\Sigma}$$

The Reduced SVD

2 of 2


In looking at the reduced SVD in this form

$$AV = \hat{U}\hat{\Sigma}$$

we note that $A \in \mathbb{C}^{m \times n}$ (if $\text{rank}(A) = n$), $V \in \mathbb{C}^{n \times n}$ (unitary), $\hat{U} \in \mathbb{C}^{m \times n}$ (unitary), and $\hat{\Sigma} \in \mathbb{R}^{n \times n}$ (diagonal, real).

If we multiply by V^* from the right, and use the fact that $VV^* = I$, we get the reduced SVD in its standard form:

$$A = \hat{U}\hat{\Sigma}V^*$$



From the Reduced to the Full SVD

In most applications the SVD is used as we have described (*i.e.* the reduced SVD “version”).

However, the SVD can be extended as follows: The columns of \hat{U} are n orthonormal vectors in \mathbb{C}^m ($m \geq n$). If $m < n$, then they do not form a basis for \mathbb{C}^m .

[LINEARLY INDEPENDENT LIST EXTENDS TO A BASIS (MATH 524, NOTES#2)]

By adding an additional $(n - m)$ orthonormal columns to \hat{U} , we get a new unitary matrix $U \in \mathbb{C}^{n \times n}$.

Further, we form the matrix Σ , by adding $(n - m)$ rows of zeros at the bottom of $\hat{\Sigma}$.

The Reduced and Full SVDs

$$A = \hat{U} \hat{\Sigma} V^*$$

$$A = U \Sigma V^*$$

We can now drop the simplifying assumption that $\text{rank}(A) = n$.

If A is rank-deficient, *i.e.* $\text{rank}(A) = r < n$, the full SVD is still appropriate; however, we only get r left singular vectors \vec{u}_k from the geometry of the hyper-ellipse.

In order to construct U , we add $(n - r)$ additional arbitrary orthonormal columns. In addition V will need $(n - r)$ additional arbitrary orthonormal columns. The matrix Σ will have r positive diagonal entries, with the remaining $(n - r)$ equal to zero.

The SVD of a Matrix: Formal Definition

Definition (Singular Value Decomposition)

Let m and n be arbitrary integers. Given $A \in \mathbb{C}^{m \times n}$, a **Singular Value Decomposition** of A is a factorization

$$A = U\Sigma V^*$$

where

$$\begin{array}{lll} U \in \mathbb{C}^{m \times m} & \text{is unitary} \\ V \in \mathbb{C}^{n \times n} & \text{is unitary} \\ \Sigma \in \mathbb{R}^{m \times n} & \text{is diagonal} \end{array}$$

The diagonal entries of Σ are non-negative, and ordered in decreasing order, *i.e.* $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p \geq 0$, where $p = \min(m, n)$.

Note: We do not require $m \geq n$. $\text{rank}(A) = r \leq \min(m, n)$.

Spheres and Hyper-ellipses

Clearly[®] if A has a SVD, *i.e.* $A = U\Sigma V^*$, then A must map the unit sphere into a hyper-ellipse:

- V^* preserves the sphere, since multiplication by a unitary matrix preserves the 2-norm. (Multiplication by V^* is a rotation + possibly a reflection).
- Multiplication by Σ stretches the sphere into a hyper-ellipse aligned with the basis.
- Multiplication by the unitary U preserves all 2-norms, and angles between vectors; hence the shape of the hyper-ellipse is preserved (albeit rotated and reflected).

If we can show that every matrix A has a SVD, then it follows that the image of the unit sphere under any linear map is a hyper-ellipse; something we stated boldly on slide 15.

[®] clearly = "Are you lost yet?" 😊

Next Time: More on the SVD

We save the proof that indeed every matrix A has a SVD for next lecture.

We also discuss the connection between the SVD and (the more familiar?) eigenvalue decomposition.

Further we make connections between the SVD and the rank, range, and null-space of A ... etc...

It takes some time to digest the SVD...

We will return to the computation of the SVD later, when we have developed a toolbox of numerical algorithms.

Homework #2

Due Date in Canvas/Gradescope

Figure out how to get your favorite piece of mathematical software (e.g. Matlab, or Python) to compute the SVD, and visualize the process/results.

Use your software (NOT “hand calculation”) to solve (pp.30–31) —

- tb-4.1, and tb-4.3

Hints:

- To get started in matlab, try **help svd**, and **help plot**.
- In Python, you likely want to
 - `import numpy`
 - and then use `numpy.linalg.svd`

There are several plotting libraries for python

- `matplotlib` is `matlabesque`
- Seaborn, Plotly, Bokeh, Altair, and Pygal are other possibilities; and there also fairly convenient plotting in pandas.
- Make sure circles look like circles, and ellipses look like ellipses.

Homework AI-Policy Spring 2024

AI-era Policies — SPRING 2024

AI-3 Documented: *Students can use AI in any manner for this assessment or deliverable, but they must provide appropriate documentation for all AI use.*

This applies to ALL MATH-543 WORK during the SPRING 2024 semester.

The goal is to leverage existing tools and resources to generate HIGH QUALITY SOLUTIONS to all assessments.

You MUST document what tools you use and HOW they were used (including prompts); AND how results were VALIDATED.

BE PREPARED to DISCUSS homework solutions and AI-strategies. **Participation in the in-class discussions will be an essential component of the grade for each assessment.**