

Numerical Optimization

Lecture Notes #28 Constrained Optimization

Peter Blomgren,
(blomgren.peter@gmail.com)

Department of Mathematics and Statistics
Dynamical Systems Group
Computational Sciences Research Center
San Diego State University
San Diego, CA 92182-7720

<http://terminus.sdsu.edu/>

Fall 2018



Outline

- 1 **Constrained Optimization**
 - KKT First Order Necessary Conditions
 - Second Order Conditions

- 2 **Some Approaches**
 - Problems and Algorithms

Introduction

We have spent a lots of effort on the Unconstrained Optimization problem, we now take a very quick look at the fundamentals of Constrained Optimization — we will quickly realize that things get quite “interesting!”

Problem 0: Constrained Optimization

$$\min_{\vec{x} \in \mathbb{R}^n} f(\vec{x}) \quad \text{subject to} \quad \begin{cases} c_i(\vec{x}) = 0, & i \in \mathcal{E} \\ c_i(\vec{x}) \geq 0, & i \in \mathcal{I} \end{cases}$$

where $i \in \mathcal{E}$ are the equality constraints, and $i \in \mathcal{I}$ the inequality constraints.

The smoothness (or lack thereof) for the objective $f(\vec{x})$ and the constraint functions $c_i(\vec{x})$ will impact the difficulty of solving the problem.



The Feasible Set

With the following definition of all allowable points:

Definition (The Feasible Set)

Let

$$\Omega = \{\vec{x} \in \mathbb{R}^n : c_i(\vec{x}) = 0 \forall i \in \mathcal{E}, \text{ and } c_i(\vec{x}) \geq 0 \forall i \in \mathcal{I}\}$$

We can rewrite the problem more compactly as

Problem 1: Constrained Optimization

$$\min_{\vec{x} \in \Omega} f(\vec{x}).$$

Our goal is to state necessary and sufficient conditions for optimality.

Local Solution

Definition (Local Solution)

A point \vec{x}^* is a local solution of Problem 1 if $\vec{x}^* \in \Omega$ and there is a neighborhood \mathcal{N} of \vec{x}^* such that $f(\vec{x}) \geq f(\vec{x}^*) \forall \vec{x} \in \mathcal{N} \cap \Omega$.

Definition (Strict Local Solution)

A point \vec{x}^* is a strict local solution of Problem 1 if $\vec{x}^* \in \Omega$ and there is a neighborhood \mathcal{N} of \vec{x}^* such that $f(\vec{x}) > f(\vec{x}^*) \forall \vec{x} \in \mathcal{N} \cap \Omega$.

Definition (Isolated Local Solution)

A point \vec{x}^* is an isolated local solution of Problem 1 if $\vec{x}^* \in \Omega$ and there is a neighborhood \mathcal{N} of \vec{x}^* such that \vec{x}^* is the only local solution in $\mathcal{N} \cap \Omega$.



Smoothness

It is usually (always?) advantageous to express constraints and objectives in as smooth a way as possible; e.g. we can replace single non-smooth conditions, like

$$\text{ns } \|\vec{x}\|_1 = |x_1| + |x_2| \leq 1$$

with several smooth constraints

$$\text{s\#1 } x_1 + x_2 \leq 1$$

$$\text{s\#2 } x_1 - x_2 \leq 1$$

$$\text{s\#3 } -x_1 + x_2 \leq 1$$

$$\text{s\#4 } -x_1 - x_2 \leq 1$$

The Active Set

Definition (Active Set)

The active set $\mathcal{A}(\vec{x})$ at any feasible \vec{x} consists of the equality constraint indices from \mathcal{E} and the indices of the inequality constraint indices from \mathcal{I} for which $c_i(\vec{x}) = 0$, *i.e.*

$$\mathcal{A}(\vec{x}) = \mathcal{E} \cup \{i \in \mathcal{I} : c_i(\vec{x}) = 0\}.$$

At a feasible point \vec{x} , the inequality constraint $i \in \mathcal{I}$ is said to be active if $c_i(\vec{x}) = 0$ and inactive if $c_i(\vec{x}) > 0$.

Linear Independence Constraint Qualification

Definition (LICQ: Linear Independence Constraint Qualification)

Given a point \vec{x} and the active set $\mathcal{A}(\vec{x})$, we say that the linear independence constraint qualification (LICQ) holds if the set of active constraint gradients

$$\{\nabla c_i(\vec{x}), i \in \mathcal{A}(\vec{x})\}$$

is linearly independent.

The Lagrangian Function

Our final building block before stating the first order conditions for optimality is the:

Definition (The Lagrangian Function, $\mathcal{L}(\vec{x}, \vec{\lambda})$)

$$\mathcal{L}(\vec{x}, \vec{\lambda}) = f(\vec{x}) - \sum_{i \in \mathcal{E} \cup \mathcal{I}} \lambda_i c_i(\vec{x})$$

The Lagrange multipliers, λ_i , are used to “pull” the solution back to the feasible set.

KKT First Order Necessary Conditions

Theorem (KKT:FONC — First Order Necessary Conditions)

Suppose that \vec{x}^* is a local solution to Problem 1, that the functions f and c_i are continuously differentiable, and that the LICQ holds at \vec{x}^* . Then there is a Lagrange multiplier vector $\vec{\lambda}^*$, with components $\lambda_i(\vec{x}^*)$, $i \in \mathcal{E} \cup \mathcal{I}$, such that the following conditions are satisfied at $(\vec{x}^*, \vec{\lambda}^*)$:

$$\begin{aligned}\nabla_{\vec{x}} \mathcal{L}(\vec{x}, \vec{\lambda}) &= 0, \\ c_i(\vec{x}^*) &= 0, \quad \forall i \in \mathcal{E} \\ c_i(\vec{x}^*) &\geq 0, \quad \forall i \in \mathcal{I} \\ \lambda_i^* &\geq 0, \quad \forall i \in \mathcal{I} \\ \lambda_i^* c_i(\vec{x}^*) &= 0, \quad i \in \mathcal{I} \cup \mathcal{E}.\end{aligned}$$

The Karush–Kuhn–Tucker conditions.

KKT First Order Necessary Conditions (compact form)

Theorem (KKT:FONC — Compact Form)

Suppose that \vec{x}^* is a local solution to Problem 1, that the functions f and c_i are continuously differentiable, and that the LICQ holds at \vec{x}^* . Then there is a Lagrange multiplier vector $\vec{\lambda}^*$, with components $\lambda_i(\vec{x}^*)$, $i \in \mathcal{E} \cup \mathcal{I}$, such that the following conditions are satisfied at $(\vec{x}^*, \vec{\lambda}^*)$:

$$0 = \nabla_{\vec{x}} \mathcal{L}(\vec{x}, \vec{\lambda}) = \nabla f(\vec{x}^*) - \sum_{i \in \mathcal{A}(\vec{x}^*)} \lambda_i^* \nabla c_i(\vec{x}^*).$$

Strict Complementarity

Definition (Strict Complementarity)

Given a local solution \vec{x}^* of Problem 1, and a vector $\vec{\lambda}^*$ satisfying the KKT:FONC, we say that the strict complementarity condition holds if exactly one of λ_i^* or $c_i(\vec{x}^*)$ is zero for each index $i \in \mathcal{I}$. In other words, we have $\lambda_i^* > 0 \forall i \in \mathcal{I} \cap \mathcal{A}(\vec{x}^*)$.

We sweep the proof of KKT:FONC under our infinitely stretchable rug. Not because it is not important (it is!), but we are somewhat short on time.

Linearized Feasible Directions

Definition (Set of Linearized Feasible Directions)

Given a feasible point \vec{x} and the active constrain set $\mathcal{A}(\vec{x})$, the set of linearized feasible directions is

$$\mathcal{F}(\vec{x}) = \left\{ \vec{d} \text{ such that } \begin{array}{l} \vec{d}^T \nabla c_i(\vec{x}) = 0, \quad \forall i \in \mathcal{E} \\ \vec{d}^T \nabla c_i(\vec{x}) \geq 0, \quad \forall i \in \mathcal{A}(\vec{x}) \cap \mathcal{I} \end{array} \right\}$$

Second Order Conditions

Second order conditions will help determine the impact of directions $\vec{w} \in \mathcal{F}(\vec{x}^*)$ for which $\vec{w}^T \nabla f(\vec{x}^*) = 0$, *i.e.* directions which are “locally flat.”

From this point on we need the functions f and c_i to be twice continuously differentiable.



Critical Cone

Given $\mathcal{F}(\vec{x}^*)$, and some Lagrange multiplier vector λ^* satisfying KKT:FONC, we define:

Definition (Critical Cone)

$$C(\vec{x}^*, \vec{\lambda}^*) = \{ \vec{w} \in \mathcal{F}(\vec{x}^*) : \nabla c_i(\vec{x}^*)^T \vec{w} = 0, \forall i \in \mathcal{A}(\vec{x}^*) \cap \mathcal{I} \text{ with } \lambda_i^* = 0 \}.$$

Or equivalently

$$\vec{w} \in C(\vec{x}^*, \vec{\lambda}^*) \Leftrightarrow \begin{cases} \nabla c_i(\vec{x}^*)^T \vec{w} = 0, & \forall i \in \mathcal{E} \\ \nabla c_i(\vec{x}^*)^T \vec{w} = 0, & \forall i \in \mathcal{A}(\vec{x}^*) \cap \mathcal{I} \text{ with } \lambda_i^* > 0 \\ \nabla c_i(\vec{x}^*)^T \vec{w} \geq 0, & \forall i \in \mathcal{A}(\vec{x}^*) \cap \mathcal{I} \text{ with } \lambda_i^* = 0 \end{cases}$$

The critical cone $C(\vec{x}^*, \vec{\lambda}^*)$ contains the directions from $\mathcal{F}(\vec{x}^*)$ for which it is not clear from first derivative information whether f will increase or decrease.

Second Order Necessary Conditions

Theorem (Second Order Necessary Conditions)

Suppose that \vec{x}^* is a local solution of Problem 1, and that the LICQ condition is satisfied. Let $\vec{\lambda}^*$ be the Lagrange multiplier vector for which the KKT:FONC are satisfied. Then

$$\vec{w}^T \nabla_{\vec{x}\vec{x}}^2 \mathcal{L}(\vec{x}, \vec{\lambda}) \vec{w} \geq 0, \quad \forall \vec{w} \in C(\vec{x}^*, \vec{\lambda}^*).$$

Interpretation: The Hessian of the Lagrangian has non-negative curvature along critical directions.

Second Order Sufficient Conditions

Theorem (Second Order Sufficient Conditions)

Suppose that for some feasible point $\vec{x}^* \in \mathbb{R}^n$ there is a Lagrange multiplier vector $\vec{\lambda}^*$ such that KKT:FONC are satisfied. Suppose also that

$$\vec{w}^T \nabla_{\vec{x}\vec{x}}^2 \mathcal{L}(\vec{x}, \vec{\lambda}) \vec{w} > 0, \quad \forall \vec{w} \in C(\vec{x}^*, \vec{\lambda}^*), \quad \vec{w} \neq \vec{0}.$$

Then \vec{x}^* is a strict local solution for Problem 1.

Much remains to be said; however, everything grows out of these fundamental definitions and theorems; leveraging special cases, weakening and strengthening conditions, and looking for alternatives.

Some Approaches

- **Linear Programming — The Simplex Method**
 - f and c_i linear functions
 - Leonid Kantorovich, 1939 — Linear Programming.
 - George Datzig, 1947 — The Simplex Method.
 - John von Neumann, 1947 — Theory of Duality.
 - The *worst case complexity* for The Simplex Method is exponential, but it is remarkably efficient in practice.

Some Approaches

- **Interior Point Methods, Primal-Dual Methods**

- c_i are *strict* inequalities.
- Better theoretical behavior than The Simplex Method.
- Leonid Khachiyan, 1979 — The Ellipsoid Method (polynomial runtime, $\mathcal{O}(n^6 L)$)
- Narendra Karmarkar, 1984 — Projective Algorithm, $\mathcal{O}(n^{3.5} L^2 \cdot \log L \cdot \log \log L)$, where n is the number of variables and L is the number of bits of input to the algorithm.



Some Approaches

• Quadratic Programming

- Special case: Exploit the structure of the problem:
- Quadratic Objective, Linear Constraints
- Active Set methods
- Interior Point methods
- Gradient Projection methods
- Appears as sub-problems for: Sequential Quadratic Programming, Penalty/Augmented Lagrangian Methods, and Interior Point Methods.

Some Approaches

- **Penalty / Augmented Lagrangian Methods**

- Constraints are represented by additions to the objective
- *Quadratic Penalty Terms* — add the square of the constraint discrepancies: intuitive, fairly simple to implement
- *Non-smooth Penalty Terms* — ℓ_1 and ℓ_0 penalty functions
- *Method of Multipliers* — estimated for the Lagrange multipliers are used.



Index